

Synthesis, Characterisation and Antibacterial activity of Metal complexes from Mannich base, N-[1-morpholino(4-nitrobenzyl)]acetamide

M.Pechiammal¹, S. Ravichandran*² and Leena Singh³

¹Department of Chemistry, Suguna College of Engineering, Nehru Nagar, Coimbatore-641 014, India

²Department of Chemistry, Veltech Dr.RR &Dr.SR Technical University, Avadi, Chennai-600 062, India

³Dept of Chemistry, Galgotia's College of Engg.&Tech., 1, Knowledge Park, Phase II, Greater Noida-201 306, India

*Corres.author: ravichanduru@yahoo.com


Abstract: Neutral complexes of Cu(II), Co(II), Ni(II) and Zn(II) have been synthesised from a new Mannich base, N-[1-morpholino(4-nitrobenzyl)]acetamide (MNBA) derived by the condensation of morpholine, 4-nitrobenzaldehyde and acetamide. The ligand forms 1:1 (metal:ligand) type of complexes with Cu(II), Co(II), Ni(II) and Zn(II) metal salts. The structural features have been arrived from their microanalytical, IR, UV-Vis., CV, EPR spectral data. The electrolytic behaviour of the chelates was assessed from their molar conductance data. The magnetic susceptibility measurements suggested that all the complexes were paramagnetic except Ni and Zn, which were diamagnetic, and the magnitude of magnetic moment values were useful to find out the number of unpaired electrons which in turn were useful to further support the geometry suggested by electronic spectral data. The magnetic susceptibility and electronic absorption spectra of copper complex indicates an octahedral geometry around the central metal ion while cobalt, zinc complexes exhibit tetrahedral geometry and nickel complex shows square-planar structure. The electrochemical behaviour, the anodic and cathodic potential and the number of electron transfer were calculated using cyclic voltammogram. The cyclic voltammogram of copper complex in MeCN solution at 298 K was studied. The X-band EPR spectra of copper complex in DMSO at 300 K and 77 K were recorded and their salient features are discussed. The antimicrobial activity of the ligand and its complexes has been extensively studied on microorganisms such as *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli* and *Pseudomonas aeruginosa* by well-diffusion technique using DMSO as solvent. The values of zone of inhibition were found out at 37°C for a period of 24 h. It has been found that all the complexes have higher activity than the free ligand and the standard.

Keywords: N-[1-morpholino(4-nitrobenzyl)]acetamide; Mannich base; antimicrobial activity.

Introduction

From the survey of existing literature, it appears that metal complexes of Mannich bases have played a vital role in the development of coordination chemistry. It is well known from the literature that the compounds containing amide moiety have a strong ability to form metal complexes and exhibit a wide range of biological activities¹⁻⁵. Earlier work reported that some drugs showed increased activity when administered as metal chelates rather than as organic compounds. Keeping the above facts in mind and as part of our continuing efforts to investigate transition metal(II) complexes using Mannich bases⁶⁻¹⁰, in this paper we describe the synthesis, characterization, redox and antimicrobial studies of Cu(II), Co(II), Ni(II) and Zn(II) complexes

containing bidentate Mannich base derived by condensing morpholine, 4-nitrobenzaldehyde and acetamide. The ligand system coordinates to the metal ion in a bidentate manner through the amide carbonyl oxygen and the nitrogen atom of morpholine group. The proposed structure of the complexes is shown below.



Experimental

All the chemicals used were of AnalaR grade. The solvents were dried and distilled before use according to standard procedures. The supporting electrolyte, tetramethylammoniumperchlorate, Me_4NClO_4 (**TMAP**) is used in the voltammetric experiment, was purchased from Sigma. IR spectra were recorded on a Jasco FT-IR-5300 instrument (KBr pellet technique). The ^1H NMR spectra were recorded at Madurai Kamaraj University, on a Bruker instrument using TMS as internal standard. The UV-Vis. spectra of all the complexes were recorded in DMSO on a Shimadzu UV-1601 spectrophotometer. Microanalytical data were performed at the Regional Sophisticated Instrumentation Centre, Central Drug Research Institute, Lucknow. Magnetic susceptibility measurements of the complexes in the solid state were determined by Gouy balance using CuSO_4 as the calibrant. Molar conductance values of the complexes were measured in DMSO at room temperature using Systronic conductivity bridge type 305. Electrochemical measurement was carried out in Electrochemical analyser model BAS-50 Voltammograph. The three-electrode cell contained a reference Ag/AgCl electrode, Pt wire auxiliary electrode and glassy carbon working electrode. EPR spectra of the copper complex were recorded on a Varian E112 X-band spectrometer at the Regional Sophisticated Instrumentation Centre, Indian Institute of Technology, Chennai using diphenylpicrylhydrazyl (**DPPH**) as the internal standard. Mueller-Hinton agar was used for testing the susceptibility of microorganisms to antibacterial agents using the well-diffusion technique. Ampicillin was used as the standard.

Synthesis of Mannich base: The Mannich base, was synthesised by the condensation of an ethanolic solution of 4-nitro benzaldehyde, morpholine and acetamide were taken in 1:1:1 mole ratio. Acetamide (5.9 g, 0.1 mol) was mixed with morpholine (10 mL, 0.1 mol) and stirred to get a clear solution. 4-nitro benzaldehyde (1.51 g, 0.1 mol) was then added in small quantities to the mixture and stirred under ice-bath condition. At first, a yellow sticky mass appeared. It was kept aside with the mother liquor open to atmosphere for *ca.* 5 days. The yellow solid formed was separated by filtration, washed with distilled water, carbon tetrachloride and recrystallised from ethanol. Yield : 60 % ; m.p : 170 $^\circ\text{C}$.

Synthesis of complexes: An ethanolic solution of Mannich base (5 mM) was mixed with metal(II) chloride (5 mM) in ethanol (20 mL) solution keeping ligand-metal ratio 1:1. The reaction mixture was then warmed for *ca.* 1 h on a water bath till the complex precipitated out. The solid complex obtained was removed by filtration, successively washed with water, dried at room temperature and recrystallised from ethanol.

Results and Discussion

All the metal complexes are stable at room temperature. They are insoluble in water but soluble in MeCN, DMF and DMSO. The ligand L, on interaction with Cu(II), Co(II), Ni(II) and Zn(II) chlorides, yields complexes corresponding to the general formula [ML]. This stoichiometric assignment is supported by the microanalytical data. The low molar conductance values of the complexes (2.6-6.8 $\text{mho cm}^2 \text{mol}^{-1}$) support their neutral nature⁸.

The FAB mass spectra of the Mannich base ligand and its copper complex are used to compare their stoichiometric composition. The Mannich base ligand shows a molecular ion peak M^+ at $m/z = 282$. The molecular ion peak for $[\text{CuCl}_2.\text{MNBA}.\text{(H}_2\text{O)}_2]$ complex was observed at $m/z = 452$, which confirms the stoichiometry of metal chelates as [ML].

In order to study the binding mode of Mannich base to metal in the complexes, IR spectrum of the free ligand was compared with the spectra of the metal complexes. Infrared spectrum of the ligand, N-[1-morpholino(4-nitrobenzyl)]acetamide and its metal complexes were recorded in KBr medium. Upon complexation with metal salts, the amide $\nu_{C=O}$ and ν_{C-N-C} of morpholine bands at 1640 cm^{-1} and 1160 cm^{-1} are shifted to lower frequencies *viz.*, $1610\text{-}1630\text{ cm}^{-1}$ and $1110\text{-}1120\text{ cm}^{-1}$ respectively. The lowering in frequencies observed in all the complexes shows the involvement of carbonyl oxygen and tertiary morpholine nitrogen atom in coordination to the metal ion. Several evidences⁷⁻¹¹ on the coordination of substituted acetamide through carbonyl oxygen have been reported.

The IR spectra of the metal complexes also show some new bands in the region $530\text{-}540\text{ cm}^{-1}$ and $440\text{-}450\text{ cm}^{-1}$ due to M-O and M-N bonds respectively which further confirm that the ligand is bidentate in nature. In all the complexes, an additional medium band found at 325 cm^{-1} is assigned to M-Cl stretching vibration. In the spectra of all the complexes, the N-H band remained at the same position as in the free ligand, indicating that the secondary nitrogen is not coordinated.

The electronic absorption spectra of the Mannich base and its Cu(II), Co(II), and Ni(II) complexes were recorded at room temperature using DMSO as solvent. The electronic spectrum of green Cu(II) complex of MNBA shows a broad band at $14,960\text{ cm}^{-1}$ assignable to ${}^2E_g \rightarrow {}^2T_{2g}$ transition, suggests an octahedral geometry for the complex. Though three transitions are expected in this case, they are very close in energy and often appear in the form of one broad band envelope. The Co(II) complex of MNBA exhibits a band with maxima at *ca.* $15,590\text{ cm}^{-1}$ which is assigned to ${}^4A_2 \rightarrow {}^4T_1(P)$ for tetrahedral geometry for the complex. The Ni(II) complex of MNBA showed band at *ca.* $16,580\text{ cm}^{-1}$ which is assigned as ${}^1A_{1g} \rightarrow {}^1A_{2g}$ transition confirming a square-planar geometry for the complex. The absence of any band below 10000 cm^{-1} eliminates the possibility of a tetrahedral environment in this complex.

The magnetic moment value of Cu(II) complex is 1.9 B.M. which suggests an octahedral geometry around the metal ion. The magnetic moment of Co(II) complex is 3.9 B.M. which suggests the high spin four coordinated tetrahedral arrangement of ligand molecules around the metal ion. The Zn(II) complex is found to be diamagnetic as expected for d^{10} configuration.

The ${}^1\text{H-NMR}$ spectrum of Mannich base ligand and its Zn(II) complex were recorded by employing TMS as internal reference at ambient temperature. The ${}^1\text{H-NMR}$ spectrum of the Mannich base ligand exhibits a multiplet signal at $\delta\ 7.2\text{-}7.8$ (m, Ar-H) and $\delta\ 5.7\text{-}5.8$ (d, N-H). In Zn(II) complex, the N-H proton is shifted slightly downfield at $\delta\ 5.8\text{-}5.9$ which reveals the bonding of the amide carbonyl oxygen to Zn(II) ion. The ${}^1\text{H-NMR}$ spectrum of the ligand show signal at $\delta\ 2.5$ for morpholine N-CH₂ and $\delta\ 3.6$ due to morpholine O-CH₂. In Zn(II) complex, the signal due to morpholine N-CH₂ protons also shifted slightly and appeared at $\delta\ 2.7$ in the complex. This is an indication of the coordination of morpholine nitrogen to the metal ion.

The EPR spectrum of copper complex provides informations which are important in studying the metal ion environment¹²⁻¹⁴. The EPR spectra of the Cu(II) complex (Figure 1) were recorded in DMSO at LNT and at RT. The spectrum of the copper complex at RT shows one intense absorption band in the high field and is isotropic due to the tumbling motion of the molecules. However, this complex at LNT shows four well resolved peaks with low field region.




Figure 1. The X-band EPR spectrum of the copper complex in DMSO at 300 K (a) and 77 K (b)

The copper complex exhibits the g_{\parallel} value of 2.32 and g_{\perp} value of 2.17. These values indicate that the Cu(II) lies predominantly in the $d_{x^2-y^2}$ orbital. The spin-orbit coupling constant, λ value (-498 cm^{-1}) calculated using the relations, $g_{av} = 1/3[g_{\parallel} + 2g_{\perp}]$ and $g_{av} = 2(1 - 2\lambda / 10Dq)$, is less than the free Cu(II) ion (-838 cm^{-1}) which also supports covalent character of M-L bond in the complex. The G value of 3.92 indicates negligible exchange interaction of Cu-Cu in the complex. The covalency parameter α^2 is calculated ($\alpha^2 = 0.85$) using the following equation:

$$\alpha^2_{cu} = -(A_{\parallel}/0.036) + (g_{\parallel} - 2.0023) + 3/7(g_{\perp} - 2.0023) + 0.04$$

If the value of $\alpha^2 = 0.5$, it indicates a complete covalent bonding, while the value of $\alpha^2 = 1.0$ suggests a complete ionic bonding. The observed value of α^2 (0.87) of the complex is less than unity which indicates that the complex has some covalent character in the ligand environment.

The cyclic voltammogram of the Cu(II) complex (0.01 M) in MeCN solution in the absence of molecular oxygen at room temperature in 1.0 to -1.2 V potential range indicates quasi-reversible one-electron process. A noteworthy feature has been observed in the cyclic voltammogram of Cu(II) complex (Figure 2).




Figure 2. Cyclic voltammogram of copper complex in MeCN

During the forward scan, it shows two cathodic reduction peaks, one at $+0.23$ V and another at -0.82 V which are attributed to reduction of $\text{Cu(II)} \rightarrow \text{Cu(I)}$ and $\text{Cu(I)} \rightarrow \text{Cu(0)}$ respectively. During the reverse scan, it shows two anodic oxidation peaks, one at -0.74 V and another at $+0.58$ V which are attributed to oxidation of $\text{Cu(0)} \rightarrow \text{Cu(I)}$ and $\text{Cu(I)} \rightarrow \text{Cu(II)}$ respectively.

Antibacterial activity of the ligand and its complexes have been carried out against the Gram positive bacteria like *S. aureus*, *B. subtilis* and Gram negative bacteria such as *E. coli*, *P. auroginosa* using Mueller-Hinton agar by well-diffusion method¹⁵ using DMF as solvent. Ampicillin was used as the standard for comparing the results.

Table 1. Antibacterial activity of ligand and its metal complexes

MNBA = Mannich base ligand

No.	Compound	Inhibition zone (mm)			
		<i>S.aureus</i>	<i>E.coli</i>	<i>P.auroginosa</i>	<i>B.subtilis</i>
1	Ligand	11	13	12	10
2	$[\text{CuCl}_2 \cdot \text{MNBA} \cdot (\text{H}_2\text{O})_2]$	16	22	21	21
3	$[\text{CoCl}_2 \cdot \text{MNBA}] (\text{H}_2\text{O})_2$	18	20	19	16
4	$[\text{NiCl}_2 \cdot \text{MNBA}] (\text{H}_2\text{O})_2$	18	21	22	19
5	$[\text{ZnCl}_2 \cdot \text{MNBA}] (\text{H}_2\text{O})_2$	20	23	21	22
6	Ampicillin	11	11	11	10

The zone of inhibition values was determined at the end of an incubation period of 24 h at 35°C . During this period, the test solution diffused and the growth of the inoculated microorganisms was affected. It has been

observed from the result (Table 1) that the metal complexes have a higher activity than that of the free ligand and the standard. Probably this may be due to the greater lipophilic nature of the complexes. Such increased activity of the metal chelates can be explained on the basis of Overtone's concept and Chelation theory.

According to Overtone's concept of cell permeability the lipid membrane that surrounds the cell favours the passage of only lipid soluble materials due to which liposolubility has important factor which controls the antimicrobial activity. On chelation, the polarity of the metal ion will be reduced to a greater extent due to the overlap of the ligand orbital and partial sharing of positive charge of metal ion with donor groups. Further, it increases the delocalisation of π -electrons over the whole chelate ring and enhances the lipophilicity of the complex. This increased lipophilicity enhances the penetration of the complexes into lipid membrane and blocking the metal binding sites on enzymes of microorganisms.

Acknowledgements

The author MP grateful to the Principal of Suguna College of Engineering, Coimbatore. Dr.SR thanks the Managing Board of Veltech University, Chennai for their encouragement. Dr.Leena thanks to Galgotia's College of Engg.& Tech., for their constant support. The authors also thank the RSIC, CDRI, Lucknow and RSIC, IIT, Chennai and Mumbai for the use of their instrumentation facilities.

References

1. Panchal P K and Patel M N, "Metal-based Antimicrobial agents: Synthesis, characterization and Biological studies of Mannich base derivatives of Benzimidazole and their metal complexes" Synth. React. Inorg. Met.-Org. Chem., 2004, 34: 1277.
2. Prabhu S, Abdul Jameel A, Ali Padusha and Syed M "Synthesis, characterization and antimicrobial activity of novel mannich bases" Der Chemica Sinica, 2012, 3(5) : 1098.
3. Sharma, Suba, Ahmad and Shamin, "Synthesis, characterization and biological activities of new mannich bases and their transition metal complexes". Journal of Chemical & Pharmaceutical Research, 2013 ,5(10) :306.
4. Muruganandam L, Krishnakumar K "Synthesis, Characterization and Antimicrobial Studies of a New Mannich Base N-[Morpholino (phenyl)methyl]acetamide and Its Cobalt(II), Nickel(II) and Copper(II) Metal Complexes". E-Journal of Chemistry, 2012, 9 (2):875.
5. Abdul Jameel, A, Palanisamy M and Syed Ali Padusha M. "Synthesis, characterization and antimicrobial studies of metal complexes of Mannich Base derived from pyridine-2-Carboxaldehyde ". Der Chemica Sinica, 2012, 3(4) 860.
6. Raman N and Ravichandran S, "Synthesis, Characterization and Antibacterial activity of a New Mannich Base, N-(1-piperidinobenzyl)benzamide and its Transition Metal(II) Complexes", Int. J. Chem.Sci., 2004, 2(2):191-198.
7. Raman N and Ravichandran S, New neutral Schiff base and its metal complexes derived from Mannich base, N-(1-morpholinobenzyl)acetamide, Polish J. Chem., 2005 ,79: 1107-1114.
8. Raman N and Ravichandran S, Synthesis and characterization of a new Schiff base and its metal complexes derived from the Mannich base, N-(1-piperidinobenzyl)acetamide, Synth. React. Inorg. Met.Org.Nano- Metal Chem., 2005, 35(6): 439 - 444.
9. Amali Roseline Emelda, Jeyachandramani N and Ravichandran S, "Synthesis, characterisation and antibacterial activity of metal complexes derived from Mannich base, N-[1-piperidinobenzyl]acetamide and 1,2- diaminobenzene", Int. J. Chem. Sci., 2007, 5(1):175-180.
10. Amali Roseline Emelda, Jeyachandramani N and Ravichandran S, "Synthesis, characterization and antimicrobial activity of Cu(II), Co(II), Ni(II) and Zn(II) complexes derived from a new Mannich base, N-(1-morpholinobenzyl)benzamide and 1,2-Diaminobenzene", Asian J.Chem., 2008, 20(1): 337-342 .
11. Nakamoto K , "Infrared and Raman Spectra of Inorganic and Coordination Compounds", 1978, 3rd Edn., John Wiley, New York.
12. Cotton F A and Wilkinson G, "Advanced Inorganic Chemistry", Wiley, New York, 1988, p.730.
13. Silverstein R M, Bassler G C and Movril T C, "Spectroscopic Identification of Organic Compounds", Wiley, New York, 4th edn., 1981, p.112.
14. Sathya D, Senthil Kumar J, Priya S, Jayachandramani N, Mahalakshmi S and Amali Roseline Emelda, "Synthesis, characterization and in vitro antimicrobial studies of some transition metal complexes with a new Mannich base N-(1-morpholinosalicylyl)acetamide", Int. J. Chem. Tech. Res, 2011, 3(1):248-252.
15. Collin C H and Lyne P M, "Microbiological methods", Butterworth, London, 3rd Edn., 1970.