



## **The effect of biotic and abiotic elicitors on Dianthalexin production from the callus of *Dianthus caryophyllus***

**Bushra Mohammed Alwash<sup>1\*</sup> and Sumaya Fadhil Hamad<sup>1</sup>**

<sup>1</sup>Department of Biology, College of Science for women, University of Baghdad/ Iraq.

**Abstract :** Biotic (*Fusariumoxysporum*) and abiotic (salicylic acid) as an elicitors were examined on the induction of phytoalexin (Dianthalexin) from carnation (*Dianthus caryophyllus*) callus from the leaf. MS medium with BA in concentrations (0.0, 0.5,1.0, 1.5, 2.0 or 2.5) mg/l and 2,4-D (0.0, 1.0, 2.0, 3.0, 4.0 or 5.0) mg/l which use for callus induction, fungalsolution in (0.0, 2.0 or 4.0) ml/l and ( 0.0, 1.0 or 2.0) mg/l of salicylic acid were added to medium. Alcoholic extraction of callus tissues was analyzed by high-performance liquid chromatography (HPLC). The results showed that the dianthalexin highest level (58.29) µg/ml on MS medium with 1.0 mg/L BA and 2,4-D and fungal elicitor was used at 1.0ml/l in three days period followed salicylic acid (58.19) µg/ml after six days incubation period.

**Keywords :** Phytoalexin, elicitors, *Fusariumoxysporum*, callus induction.

**Bushra Mohammed Alwash *et al*** /International Journal of ChemTech Research, 2017,10(6): 25-30.

\*\*\*\*\*