



## **Synthesis and Characterization of Bis-Pyrazoles Using 1-(5-Methyl Pyridine-2-yl) Pyrrolidine-2,5Dione Under Microwave Irradiation**

**S.S.Rajput<sup>1\*</sup>, S.N.Patel<sup>2</sup>, S.B.Chaudhari<sup>3</sup>**

<sup>1\*</sup> Department of Chemistry, SVS's DadasahebRawal College Dondaicha 425408, Maharashtra, India

<sup>2</sup> Department of Chemistry, S.P.D.M. College, Shirpur 425405, Maharashtra, India

<sup>3</sup> Departments of Chemistry, R.C.Patel ASC College, Shirpur 425405, Maharashtra, India

**Abstract:**The current study shows that the green and efficient synthesis of new series of bis-pyrazoles derivatives from 1-(5-methyl pyridine-2-yl) pyrrolidine-2,5dione under microwave irradiation. The 1-(5-Methylpyridin-2-yl) pyrrolidine-2,5-dione, was synthesized from succinic anhydride and 5-methyl pyridine-2-amine. This cyclic imides condensed with substituted aromatic aldehydes using solid support neutral  $Al_2O_3$  in microwave. The resulting bis-chalcone derivatives underwent ring closer with hydrazine hydrate in presence of neutral  $Al_2O_3$  under microwave irradiation offered bis-pyrazoles derivatives.

**Keywords:** Succinic anhydride, 5-methyl pyridine-2-amine, bis-chalcone, bis-pyrazole.

**S.S.Rajput et al**/International Journal of ChemTech Research, 2017,10(9): 73-78.

\*\*\*\*\*