

Antifertility Effect And Hormonal Profile of Petroleum Ether Extract of Seeds of *Cassia fistula* in Female Rats

Rajesh Yadav^{1*} and G. C. Jain²

¹ Department of Zoology , Mahatma Gandhi Institute Of Applied Sciences, Jaipur (India)

² Reproductive Physiology Laboratory, Centre for Advanced Studies in Zoology,
University of Rajasthan, Jaipur - 302004 (India)

*E-mail: ryadav70@yahoo.co.in

Abstract : Petroleum ether extract of seeds of *Cassia fistula* was screened for the antifertility activity in proven fertile female albino rats at the doses 100, 200 and 500mg/kg b.wt./day. Oral administration of the extract to mated female rats on days 1-5 of pregnancy resulted in a decline in the fertility index, numbers of uterine implants and live fetuses in a dose dependent manner as was confirmed by laparotomy on day 15 of pregnancy. The extract (100mg/kg b.wt.) exhibited weak estrogenic activity when given alone and tested in immature bilaterally ovariectomized female albino rats, but exhibited slight antiestrogenic activity when administration along with estradiol valerate (0.1mg/kg b.wt.). Blood sugar and haematological parameters were within normal range. Thus, the results of the present study indicate that the petroleum ether extract of *Cassia fistula* seeds possesses pregnancy terminating effect by virtue of anti-implantation activity.

Key words : *Cassia fistula*, antifertility, antiimplantation, female rats

Introduction

The quest for the oral contraceptive agent that can control human fertility is as old as recorded history. Although a variety of synthetic contraceptive agents are available, but these can not be used continuously because of their side effects. So, natural plant substances possessing mild inherent estrogenic and antiestrogenic properties offer themselves as an effective non-conventional source of contraception with less deleterious side effects.

Cassia fistula Linn. (Hindi-Amaltas; English-Golden Shower or Indian Laburnum), a medium sized tree belonging to the family - Caesalpiniaceae, is widely cultivated throughout India as an ornamental tree. *Cassia fistula* has been used extensively in the folklore medicine for the treatment of a variety of diseases¹⁻². Pharmacologically the plant has been investigated for its antibacterial³, anti-diabetic⁴, hypocholesterolaemic⁵, hepatoprotective⁶, antitumour⁷, laxative⁸ and antioxidant⁹ effects. The plant is rich in phenolic antioxidants such as anthraquinones, flavonoids and flavan-3-ol derivatives¹⁰. Ethanolic extract of fruits of this plant has been reported to possess anti-implantation and estrogenic effect in rats¹¹. The previous study in our laboratory showed that postcoital administration of aqueous extract of seeds of

Cassia fistula at the dose 500mg/kg b.wt./day prevented pregnancy in all the treated female rats by virtue of anti-implantation activity¹². Because of the wide range of therapeutic efficacies of *Cassia fistula* plant and since the active principles present in various extracts of it may be different, the present study, therefore, proposes to evaluate the post-coital antifertility efficacy of petroleum ether extract of seeds of *Cassia fistula* in female rats and also to investigate its hormonal profile in immature bilaterally ovariectomized female rats in order to gain insight into its possible mechanism of action.

Materials and Methods

Plant material and extraction

Cassia fistula pods were collected during the season and were thoroughly dried in the shade. The plant was authenticated at the Department of Botany, University of Rajasthan, Jaipur (India). The seeds separated from the shade dried pods were ground to coarse powder (1000g) and was subjected to soxhlet extraction with petroleum ether (B.P.60°-80°). The crude extract thus obtained was concentrated under reduced pressure and low temperature. The residue obtained was then utilized for evaluating antifertility efficacy by

suspending in appropriate volume of olive oil. The yield of the extract was 12% of starting crude material.

Experimental animal

Colony bred, adult albino Wistar rats (weighing 170-200 g) for antifertility studies and immature female rats (21-24 days old) for bioassay studies were used as experimental animal model. All the animals were housed in standard laboratory conditions (temperature $22 \pm 3^\circ\text{C}$ and 14hr light/10hr dark cycle) with free access of food (Lipton India Ltd) and tap water *ad libitum*. All the experimental procedures were performed according to the guidelines for the care and use of experimental animals and approved by the Institutional Ethical Committee for Animals Care and Use, University of Rajasthan, Jaipur (India).

Dose and route of administration

The animals of Group I received vehicle (olive oil, 0.2ml/rat) only and served as control. Animals of Group II, III and IV received crude petroleum ether extract of *Cassia fistula* at 100, 200 and 500 mg/kg b.wt./day (suspended in olive oil) doses, respectively, once a day from day 1-5 *post coitum* (*pc*). The extract was administered orally by using a curved needle and a tuberculin syringe.

Antifertility study:

For the antifertility study, only normal cycling proestrous or estrous female rats were caged over-night with males (2:1 ratio) of proven fertility. Next morning, insemination was confirmed by the presence of the vaginal plug and spermatozoa in the vaginal smear. This day of mating was designated as day 'Zero' of pregnancy. These mated females were isolated, weighed and divided into four groups of seven animals each. In order to confirm if implantation occurred following mating, all the control and treated female rats were sacrificed on day 15 *pc* under mild ether anaesthesia and their body weights were recorded. Blood samples for hematological studies were collected directly from the cardiac puncture¹³. During autopsy, both the uterine horns were examined for the number of implantation sites, live or dead / resorbed fetuses. Embryos with bright reddish aspect and clear margins were considered to be normal and those with dull blue colour, no clear margin, smaller in size and with some surrounding exudate were considered to be resorbing. The ovaries were excised and examined for the number of fresh corpora lutea using a stereomicroscope. The uterine horns were removed and trimmed of fat. These uterine horns with embryonic contents intact were quickly weighed on an electric pan-balance to the nearest milligrams. The fetuses were removed from the uterine horns and suitable parts of these uterine horns were fixed in Bouin's fixative for histological observations in future.

Hematology

The counts of RBC and WBC, Hemoglobin and hematocrit values were determined from the blood collected directly from the heart of rats receiving 500mg/kg b.wt. extract at the time of sacrifice¹³.

Hormonal profile / Estrogenic and antiestrogenic activity

Crude petroleum ether extract of the test substance was subjected to standard bioassay procedures for assessment of estrogenic or antiestrogenic activity in terms of the rat uterotrophic assay¹⁴. Immature female rats (21-24 days old) were taken for bioassay studies. These animals were bilaterally ovariectomized by dorsolateral approach under light ether anaesthesia and semisterile conditions and after a rest period of seven days, these were randomly divided into four groups of five rats each and treated as follows:

Group I : Control group, receiving olive oil only (0.2ml/rat/day), orally.

Group II : Estradiol valerate (0.1 mg/kg b.wt./day), intramuscularly (i.m.).

Group III : Extract alone (100 mg/kg b.wt./day), orally.

Group IV : Extract (100 mg/kg b.wt./day, orally) + Estradiol valerate (0.1 mg/kg b.wt./day, i.m.), conjointly.

All these rats received treatment twice daily for 3 consecutive days. These treated rats were sacrificed 24 hours after the last dose administration. Their body weights were recorded. Uteri were carefully dissected out, freed from adherent tissues, blotted on filter paper and were weighed quickly to nearest milligrams on digital balance. Condition of vaginal opening was also recorded.

Uterine luminal epithelial cell height

Haematoxylin-eosin stained slides were observed microscopically for luminal epithelial cell height. One hundred luminal epithelial cells from 25 sections were measured with ocular micrometer at X400. Two diagonal and one median lengths were measured, averaged and expressed as mean epithelial cell height and were then calibrated with a stage micrometer.

Statistical analysis

All the values are expressed as the mean \pm SEM for 7 rats per group. Data were analysed statistically by Students 't' test and $p < 0.05$ was considered as statistically significant.

Results

Antifertility activity

Table I summarizes the data obtained in the fertility study after postcoital administration of petroleum ether extract of seeds of *Cassia fistula*. In the control group, all the mated female rats were pregnant. Oral administration of petroleum ether extract of seeds of *Cassia fistula* at the doses 100, 200 & 500mg/kg b.wt. to the female rats from day 1-5 *pc* impaired fertility substantially in terms of quantal pregnancy and number of uterine implants. The quantal pregnancy rate in rats receiving 100 and 200 mg/kg b.wt./day extract doses was declined to 42.86% and 28.57%, respectively. However, the dose (500mg/kg b.wt./day) of the extract also exhibited only 28.57% quantal pregnancy as two of the females showed the presence of implantation sites on day 15 *pc*. The number of total uterine implantation sites and viable fetuses showed a dose-dependent decrease by virtue of increase in the percentage of the pre-implantation embryonic loss rate. The total number of healthy corpora lutea in control and extract treated rats remained significantly unchanged.

Ponderal changes (Body and Uterine weight) : Administration of crude petroleum ether extract of seeds of *Cassia fistula* orally at different doses from day 1-5 pc did not produce any significant change in the maternal body weight, but did produce a significant decline in relative uterine weight when compared with controls (Table 2).

Hematology

A statistically non-significant change in the RBC and WBC counts, hemoglobin and hematocrit values was observed (data not shown).

Hormonal profile : Table 3 shows the results of uterine bioassay studies of petroleum ether extract of seeds of *Cassia fistula* in bilaterally ovariectomized immature rats. Oral administration of the extract (100mg/kg b.wt./twice daily) to ovariectomized immature female rats produced a slightly significant ($p < 0.05$) increase in the uterine wet weight. However, the extract did not induce premature opening of the vagina, thus, suggesting mild estrogenic activity of the extract. But when the extract was administered conjointly with estradiol valerate (EDV, 0.1mg/kg b.wt./twice daily), it significantly ($p < 0.001$) prevented the estrogen induced uterotrophic effect, thus, reflecting antiestrogenic nature of the extract in presence of a strong estrogen.

Uterine luminal epithelial cell height : The extract when administered alone to ovariectomized rats induced a mild stimulation of all the uterine constituent elements and a significant ($p < 0.01$) increase in the uterine luminal epithelial cell height when compared with ovariectomized control only, thus, showing the estrogenic nature. But when the extract was administered along with a strong estrogen i.e. estradiol valerate (EDV), it showed antagonism of EDV induced hypertrophy of the uterine constituent elements and a highly significant ($p < 0.001$) decline in the uterine luminal epithelial cell height in comparison to EDV alone treated rats (Group II).

Discussion

In the present study oral administration of petroleum ether extract of *Cassia fistula* seeds at the doses 100, 200 and 500mg/kg orally from day 1 to 5 pc, produced a dose dependent adverse effect on fertility index (quantal pregnancy) and number of implantations in uterine horns of the female rats by virtue of an increase in the percentage of the pre-implantation embryonic loss. These results are in agreement with the earlier findings of Bhardwaj and Mathur¹¹.

The antifertility effect of aqueous extract of *Cassia fistula* has been reported earlier by our laboratory¹². The present findings indicate that petroleum ether extract of seeds of *Cassia fistula* also possess significant antifertility activity as it interfered with steroidal conditioning of the uterus and renders it hostile to ovum implantation¹⁵.

A significant decrease in relative uterine weight after postcoital administration of the extract was observed on day 15 pc in comparison to control. This decrease in uterine weight was correlated with decrease in the

number of implantation sites and viable fetuses in the uterine horns¹⁶⁻¹⁷. As, the uterine weight in pregnant rats also serves as an index of uterine decidualization and a significant decrease in uterine weight indicates suppression of decidual changes in uterus¹⁸.

In the present investigation, a non-significant change in the total erythrocyte and leucocyte counts, hemoglobin and hematocrit values following oral therapy of petroleum ether extract of seeds of *Cassia fistula* suggests non-toxic action of the extract on general body metabolism. These results are in agreement with the reports of Mutreja *et al*¹⁹ who also reported a non-significant change in haematological parameters after administration of alcoholic extract of seeds of *Nelumbo nucifera*.

In bioassay test, crude petroleum ether extract of *Cassia fistula* seeds showed mild estrogenicity when treated alone but when treated conjointly with estradiol valerate it produced estrogen antagonistic effects. Thus, the pregnancy interceptory effect of the extract of the test plant might be due to its antiestrogenic nature. Thus, like other relatively weak estrogenic substances it behaved as an antiestrogen in the presence of relatively more potent estrogen by possibly affecting the uterine estrogen receptor binding. A number of plants possessing antiestrogenic activity have also been reported to interrupt pregnancy²⁰⁻²³.

Antiestrogens with weak estrogenic activity administered early in pregnancy may interfere with implantation by altering the normal pattern of hormonal conditioning of uterus required for conception²⁴. Furthermore, many of the synthetic non-steroidal compounds having estrogen agonist-antagonistic property have also shown postcoital antifertility effects²⁵⁻²⁶. Experimental studies also indicate pregnancy blocking effect of certain synthetic estrogen agonist-antagonist compounds²⁷⁻²⁸. Pre-implantation losses can arise due to disruption of events which are pre-requisite for fertilization or an impairment in the production of cytokines, growth factor and various types of adhesion molecules either by the developing blastocyst or by the uterine epithelium around the site of implantation²⁹⁻³⁰. Therefore, one possible explanation of anti-implantation effect of the extract can be explained by pre-implantation embryonic loss due to accelerated embryonic transport which is an estrogen mediated process.

In conclusion, the present study suggests that the antifertility activity of the petroleum ether extract of seeds of *Cassia fistula* is probably due to its anti-oestrogenic property. The action of estrogen on the uterus, which is essential for implantation, is antagonized by the extract. Hence, an unfavourable environment is created in the endometrium checking nidation. While the hematological studies performed in extract treated rats did not reflect any adverse effect.

TABLE 1: EFFECT OF PETROLEUM ETHER EXTRACT OF SEEDS OF *Cassia fistula* ON IMPLANTATIONS IN FEMALE RATS FROM DAY 1-5 POST-COITUM

Group	Treatment dose (mg/Kg b.wt.)	No. of pregnant rats (Fertility index) ¹	No. of implantation sites in individual rats	No. of reabsorbing fetuses	No. of viable/live fetuses	No. of corpora lutea	%Preimplantation loss ²	Postimplantation or Reabsorption rate ³	Birth rate ⁵
Control (0.2ml olive oil/rat)	-	7 (100)	(10,11,13,13,12,13,12)	0	84	105	20	0	80
Pet. ether extract	100	3 (42.86)	(0,10,0,4,0,13, 0)	12	15	101	73.27	44.44	14.85
	200	2 (28.57)	(0,10,0, 0,12, 0, 0)	10	12	100	78	45.45	12
	500	2 (28.57)	(0,0,8,0,0,10, 0)	9	9	102	82.35	50	8.82

¹Quantal pregnancy or fertility index (%) = (No. of pregnant animals/No. of mated animals) x 100

²Preimplantation loss (%) = [(Total no. of corpora lutea – Total no. of implantation)/Total no. of corpora lutea] x 100

³Postimplantation loss (%) or Reabsorption rate(%) = [(Total no. of implantation – Total no. of viable fetuses)/Total no. of implantation] x 100

⁴Birth rate = no. of live fetuses/ no. of corpora lutea x 100

TABLE 2 : EFFECT OF PETROLEUM ETHER EXTRACT OF SEEDS OF *Cassia fistula* ON THE BODY AND RELATIVE UTERINE WEIGHT OF FEMALE RATS

Group	Treatment dose (mg/Kg b. wt.)	Initial body wt. (g)	Final body wt. (g)	uterine weight (g/100 g b. wt.)
Control (0.2ml olive oil/rat)	-	198 ± 5.83	215 ± 12.65	6.45 ± 0.27
Experimental (Pet. Ether extract)	100	174 ± 17.49	182 ± 15.62	2.52 ± 1.35*
	200	170 ± 13.04	176 ± 7.48	2.51 ± 1.47*
	500	180 ± 7.07	184 ± 5.09	2.47 ± 1.34*

[Values are mean + SEM]

Level of Significance when compared with control rats :

* p < 0.05

TABLE 3 : SHOWING ESTROGENIC AND ANTIESTROGENIC ACTIVITY OF PETROLEUM ETHER EXTRACT OF SEEDS OF *Cassia fistula* IN BILATERALLY OVARIECTOMIZED IMMATURE FEMALE RATS

Group	Treatment group	uterine weight (mg/100g b. wt.)	vaginal opening	Luminal epithelial cell height (μm)
I.	Control (0.2ml olive oil/rat)	38.36 \pm 1.63	closed	10.45 \pm 0.32
II.	Estradiol valerate (0.1mg/kg b.wt.)	529.36 \pm 23.91 ^{aaa}	open	42.50 \pm 0.29 ^{aaa}
III.	Pet. ether extract (100mg/kg b.wt.)	45.50 \pm 1.41*	closed	16.72 \pm 0.37 ^{***}
IV.	Pet. Ether extract (100mg/kg b.wt.) + Estradiol valerate (0.1mg/kg b.wt.)	308.50 \pm 13.99 ^{###}	open	35.64 \pm 0.24 ^{###}


[Values are mean \pm SEM]



1. Group II and group III treated rats compared with Group I rats.

aaa $p < 0.001$, * $p < 0.05$, *** $p < 0.001$

3. Group IV treated rats compared with Group II rats.

$p < 0.001$ (Highly significant)





References

1. Chopra RN, Nayar SL, Chopra IC. Glossary of Indian medicinal plants, Publication and information Directorate, CSIR, New Delhi. 1992: 54.
2. Barthakur NN, Arnold NP, Alli I. The Indian laburnum (*Cassia fistula* L.) fruit: an analysis of its chemical constituents. *Plants Food Hum Nutr.* 1995; 47:55-62.
3. Perumal Samy R, Ignacimuthu S, Sen A. Screening of 34 Indian medicinal plants for antibacterial properties. *J Ethnopharmacol.* 1998; 62:173-182.

4. Esposito Avella M, Diaz A, deGracia I, de Tello R, Gupta MP. Evaluation of traditional medicine: effects of *Cajanus cajan* L and of *Cassia fistula* L on carbohydrate metabolism in mice. Rev Med Panama. 1991;16:39-45.
5. El-Saadany SS, el-Massry RA, Labib SM, Sitohy MZ. The biochemical role and hypocholesterolaemic potential of the legume *Cassia fistula* in hypercholesterolaemic rats. Nahrung. 1991; 35:807-815.
6. Bhakta T, Banerjee S, Subhash C. Hepatoprotective activity of *Cassia fistula* leaf extract. Phytomedicine. 2001; 8 (3) : 220-224.
7. Gupta M, Mazumder UK, Rath N, Mukhopadhyay DK, Antitumour activity of methanolic extract of *Cassia fistula* L. seeds against Ehrlich ascites carcinoma. J Ethnopharmacol. 2000. 72: 151-156.
8. Akanmu M A, Iwalewa E O, Elujoba A A, Adelusola K A. Toxicity potentials of *Cassia fistula* fruits as laxative with reference to Senna. African J of Biomedical Research; 2004; 7(1): 23-26.
9. Luximon- Ramma A, Bahorun T , Soobrattee MA, Aruoma OI. Antioxidant activities of phenolic , proanthocyanidins and flavonoid components in extracts of *Cassia fistula*. J Agric Food Chem 2002. 50:5042-5047.
10. Bahorun T, Neergheen VS, Aruoma OI. Phytochemical constituents of *Cassia fistula*. African J Biotechnology, 15, 4 : 1530-1540.
11. Bhardwaj S, Mathur R. Antifertility screening of fruits of *Cassia fistula* in female albino rats. Comp Physiol Ecol. 1979; 4:277-279.
12. Yadav R, Jain GC. Antifertility effect of aqueous extract of seeds of *Cassia fistula* in female rats. Advances in Contraception, 1999; 15:292-301.
13. Burhoe SO. Method of securing blood from rats. J Heredity, 1940, 31: 445-449.
14. Dorfman RI, Kincl FA. Uterotrophic activity of various phenolic steroids. Acta Endocr (Kobenhavn) 1966; 52:619.
15. Wakuda K, Takakura K, Nakanishi K, Kita N, Hirose M and Noda Y. Embryo-dependent induction of embryo receptivity in the mouse endometrium. J. Reprod Fert. 1999; 115 315-324
16. Goonasekera MM, Gunawardana VK, Jayasena K, Mohammed SG and Balasubramaniam S. Pregnancy terminating effect of *Jatropha curcas* in rats. J Ethnopharmacol. 1995; 47 117-123
17. Tafesse G, Mekonnen Y, Makonnen E. Antifertility effect of aqueous and ethanol extracts of the leaves and roots of *Asparagus africanus* in rats. African Health Sci. 2006, 6: 81-85.
18. Ema M, Emiko M and Kunio K. Suppression of uterine decidualization as a cause of implantation failure induced by triphenyltin chloride in rats. Arch Toxicol. , 1999; 73: 175-179.
19. Mutreja A, Agarwal M, Kushwaha S, Chauhan A. Effect of *Nelumbo nucifera* seeds on the reproductive organs of female rats. Iranian J Of Repro Med 2008; 6(1):7-11.
20. Shukla S, Mathur R, Prakash AO. Antifertility profile of aqueous extract of *Moringa oleifera* roots. J Ethnopharmacol. 1988; 22:51-62.
21. Badami S, Aneesh R, Sankar S, Sathishkumar M N, Suresh B, Rajan S. Antifertility activity of *Derris brevipes* variety coriacea. J Ethnopharmacol. 2003; 84 (1) : 99-104.
22. Sharma JD, Bhinda Anuja. Antifertility activity of steroidal extract of *Trigonella foenum graecum* (seeds) in female rats. Asian J Exp Sci. 2005, 19(1) : 115-120
23. Ravichandran V, Suresh B, Satishkumar MN, Elango K, Srinivasan R . Antifertility activity of hydroalcoholic extract of *Ailanthus excelsa* (Roxb) : An ethnomedicines used by tribals of Nilgiris region in Tamilnadu. Ethnopharmacology, 2007, 112: 189-191.
24. Psychoyos A, Prepas I. Inhibition of egg development and implantation in rats after post-coital administration of the progesterone antagonist RU 486. J reprod fertil. 1987; 80:487-491.
25. Mehrotra pK, Shukla R, Dwivedi A, Srivastava RP, Bhat N, Seth M, Bhaduri AP, Kamboj VP. Pregnancy interceptive efficacy and biological profile of 3-amino-6,7-dimethoxy-1H-pyrazolo[3,4-b]quonoline (compound 85/83) in rodents. Contraception, 1991. 43: 507-519.
26. Singh MM, Sreenivasulu S , Kamboj VP, Duration of antiimplantation action of the triphenylethylene anti-estrogen centchroman in adult female rats. J Reprod Fert, 1994. 100: 367-374.
27. Kaunitz AM, Illions ED, Jones JL, Sang LA. Contraception : A clinical review for the Internist. Med J Clinics of North America 1995. 79: 1377-1405.
28. Dao B, Vanage G, Marshall A, Bardin CW, Koide SS. Antiimplantation activity of antiestrogens and mifepristone. Contraception 1996, 54: 253-258.
29. Denker H W. Implantation : a cell biological paradox. J Exp Zool. 1993; 266 : 541-558.
30. Haimovici F, Anderson D J. Cytokines and growth factors in implantation. Microscopy Res Tech. 1993; 25 : 201-207.
