

Physicochemical and Bioactives Characteristics of Purple and Yellow Water Yam (*Dioscorea alata*) Tubers

Harijono, Teti Estiasih, Mulia W. Apriliyanti*,
Asmak Afriliana**, Joni Kusnadi

Food Science and Technology Department, Brawijaya University, Malang,
Indonesia 65145.

Corres.author: mulia_apriliyanti@yahoo.com*, asmak_1118@yahoo.com**

Phone and Fax +62-341-56921.

Abstract: This study aimed to characterize the physico-chemical properties and identify the bioactive compounds including diosgenin, water soluble polysaccharides, and water soluble proteins containing dioscorin from purple water yam (*Dioscorea alata* var *purpurea*) and yellow water yam (*Dioscorea alata* L.). Both varieties of water yam contain water soluble polysaccharides that had CH₃, CH₂, OH, NH, C=O, acetyl (C-O), carboxyl (COOH), and C-O-C groups. WSP hydrolyzate contained more glucose. Mannose, arabinose, glucuronic acid and galacturonic acid in small quantities. Purple and yellow water yam contained 25.94 and 25.45% dioscorin from water soluble protein. The analysis of amino acid profile showed that the water soluble protein of water yam comprises of aspartate, glutamate, serine, histidine, glycine, arginine, alanine, tyrosine, valine, phenylalanine, isoleucine, leucine, and lysine. Both tubers also had diosgenin, a steroidal saponin of yam family.

Keywords: water yam, diosgenin, water soluble polysaccharides, water soluble proteins, dioscorin.

Introduction

Dioscorea family or yam tuber consists of about 600 species which are around 50-60 species cultivated for food and medicine. The well known yam family in Indonesia is *D. alata* L, *D. hispida*, *D. bulbifera* L, and *D. esculenta* that have productivity of 60-70 ton/ha, but these tubers have not yet been used intensively. *Dioscorea alata* L or water yam is a creep and shrub plant, generally cultivated in between forest plants¹. Water yam tuber is differentiated by its flesh color, namely purple, yellow, and white. It is reported that *Dioscorea* contains several bioactive compounds such as diosgenin, water soluble polysaccharides, and dioscorin^{2,3}.

Diosgenin is a phytochemical compound to prevent colon cancer and decrease cholesterol absorption^{4,5}. Water soluble polysaccharides of yam could be used to reduce the blood glucose and cholesterol levels (especially the low density lipoprotein cholesterol)^{6,7,8}. Dioscorin of *Dioscorea* tubers is a protein that binds strongly to water soluble polysaccharides⁹. *Dioscorea alata* cv. Tainong number 1 from Taiwan contains dioscorin up to 90% of water soluble protein. Dioscorin is a storage protein in some species of yam that has function as trypsin inhibitor, carbonic anhydrase, antioxidant, immunomodulator, and antihypertension^{10,11,3,12}.

The purpose of this study is to characterize the physico-chemical properties and identify bioactive compounds of purple and yellow water yams, including diosgenin, water soluble polysaccharides, and dioscorin.

Materials And Methods

Materials

One of purple (*Dioscorea alata* var *Purpurea*) and yellow water yam (*Dioscorea alata* Linn.) were obtained from a local market in Tuban, East Java, Indonesia. Reagents used for analysis and bioactive compound extraction were 96% ethanol (pa), distilled water, sulfuric acid, chloroform, trichloroacetic acid 10%, KBr, Ba(OH)₂, glucose (sigma), mannose (sigma), galactose (sigma), arabinose (sigma), rhamnose (sigma), glucuronic acid (sigma), galacturonic acid (sigma), acetonitrile, water for HPLC, Tris, HCl, acrylamide, bis-acrylamide, glycine, TEMED (N-tetramethyl-ethylenediamine), protein marker (sigma, St Louis, MO, USA), coomassie blue R-250, Na₂ EDTA, KCl, PVP (polyvinylpyrrolidone), DTT (dithiothreitol), APS (ammonium persulfat), acetate acid glacial, methanol, HCl (pa), NaOH, sodium acetate, THF (tetra hidrofuran), OPA (orthophalaldehyd), and H₂SO₄ 0.005 M.

Methods

Extraction of diosgenin

The extraction of diosgenin¹³, 10 grams of water yam tuber were cut into small pieces. Then, they were added 100 mL of ethanol-sulfuric acid 2.5M. The mixture were refluxed for 4 hours at 80°C. The solution were cooled and filtered through a Whatman filter No. 1. The filtrate were diluted to 200 mL with distilled water. The solution were extracted with chloroform 10 mL for 3 times. Chloroform extract was evaporated under 60°C. The evaporated were redissolved in chloroform to 10 mL.

Extraction of water soluble polysaccharides (WSP)

Fresh water yam tubers were peeled, cut, weighed and added distilled water with a ratio 1:3 (w/v) then they were blended and filtered. Water yam juice were centrifuged at 4500 rpm for 20 minutes. After that, supernatant were taken and precipitated with 96% ethanol (1:4) for 36 hours¹⁴. Wet water soluble polysaccharide extracts is dried at 50°C for 18 hours.

Extraction of water soluble protein

Supernatant of water yam juice were added with a ratio 1:1 of 10% trichloroacetic acid, mixed evenly, and centrifuged at 3500 rpm for 20 minutes. The obtained precipitate was washed by distilled water until neutral pH. This is modification from protein extract method biuret¹⁵.

Physico-chemical properties

Purple and yellow water yam were characterized for their physical properties by descriptive test with scoring¹⁶. Physical analysis were based on the test scoring of 20 panelists on tuber peel color, tuber shape, tuber peel texture, the abundance of mucilage, the intensity hair bulbs, color of the flesh, and texture of the flesh. The chemical analysis were moisture, ash, protein, and fat, crude fiber¹⁶, soluble fiber, insoluble fiber¹⁷, and antioxidant activity¹⁸. The analysis of soluble and insoluble fiber used the multi enzyme method.

Identification of bioactive compound

The diosgenin extract was dissolved in 1 mL of 96% ethanol and then injected on HPLC (Shimadzu), a Sunfire®C18 column (150 mL x 4.6 mm, 5 µm), the experimental conditions were an isocratic binary system of acetonitrile: water (90:10), a flow rate of 1 mL/min and a temperature of 35°C. Detection was performed at 194 nm¹⁹.

Functional groups of dry water soluble polysaccharide were analyzed by Fourier Transformation Infra Red Spectrometer (Shimadzu, model 8400S). The analysis of sugar constituent were performed by HPLC. Before

HPLC analysis, water soluble polysaccharides was hydrolyzed by strong acid²⁰. HPLC (KNAUER) with Refractive Index S2300 detector, Aminex HPX-87H column (300 x 7.8 mm), and H₂SO₄ 0.005 M as eluent.

Soluble protein extracts containing dioscorin were analyzed in Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (Bio-Rad, Richmond, USA) to identify the composition of protein based on molecular weight, and gel documentation (Chemidoc XRS, Bio rad Laboratories) were used for quantitative analysis of separated protein. Amino acids composition of water soluble protein were analyzed by HPLC (Shimadzu LC10). Licospher®100 RP 18 column (5mm), length 125 mL x 4 mm with mobile phase A were CH₃OH: 50 mM sodium acetate: THF (2:96:2) pH 6.8 and mobile phase B 65% CH₃OH (modification from AOAC, 2005), and detector of Fluorescence (Shimadzu RF-138).

Results and Discussion

Physical-chemical properties

There are variations of tuber shape regularity on various sizes of purple water yam tubers, while the yellow one was almost similar. Purple water yam peel texture more rough than yellow one. The leather peel texture of purple yam were more rough on a larger tuber. Mucus from purple water yam tuber were influenced by the tuber size, where bigger tuber has more mucus. The mucus of yellow water yams were vary and did not depend on tuber size. Yellow water yams were more hairy than purple yam. The hair were influenced by the tuber size, where the bigger tuber has more hair. The yellow water yam texture not be affected by tuber size while purple yam affected. The larger size of the tuber had more delicate texture (Table 1), which may related to the size of the tuber starch granules.


Chemical characteristics of purple water yam were similar with yellow water yam tuber (Table 2), except the water content and carbohydrate. Yellow water yam had a higher moisture content. Yellow water yam did not have economic value because of the high moisture content. Mean while, there were several sub-types of *Dioscorea alata* which had low moisture content and high level of starch²¹. Protein of water yam tuber were lower than expectation. Dioscorin was a reserve protein in Dioscorea group that medicinal properties. Likewise, the fat content were quite low, so the most contribution to energy of water yam tuber comes from carbohydrates. Crude fiber content of water yams were lower than fruits and vegetables. Soluble fiber of water yam were almost similar. Dietary fiber were often underestimated, compared with protein, vitamins, and other nutrients. Although not classified as nutrients, dietary fiber proved to be very beneficial for health. From the analysis of insoluble fiber also indicates that the water yams contain high insoluble fiber.

Table 1: The difference in physical properties of purple and yellow water yam tubers

Physical Property	Purple water yam	Yellow water yam
Color of the peel	dark brown	light brown
Tuber shape	Slightly irregular (oval and round)	Irregular (oval, round, and fingered)
Texture of the peel	rough	rather smooth
Many little mucus	many mucus	rather mucus
Presence or absence of hair	little hair	rather hair
Color of the flesh	purple	light yellow
Texture of the flesh	smooth	rather smooth

Table 2: Chemical composition of purple and yellow water yam

Component (%)	Purple water yam	Yellow water yam
Water	78.12	87.75
Ash	0.69	0.81
Protein	1.69	1.58
Fat	0.97	0.47
Carbohydrat by difference	18.53	9.39
Crude fiber	1.46	1.17
Soluble fiber	3.23	3.28
Insolublefiber	36.06	51.45

Figure 1: Chromatogram of extract diosgenin from purple (a) and yellow water yams (b)


Diosgenin

Diosgenin was a steroidal saponin from acid, base, or enzyme hydrolysis of saponins which has chemical formula of $C_{27}H_{42}O_3$. The purple and yellow water yams contained diosgenin of 0.015 g/kg and 0.006 g/kg. *Dioscorea alata* var *Purpurea* from Mumbai contained 0.78 g/kg of diosgenin². The highest content of diosgenin glycoside were in subterranean organs such as rhizomes or tubers and the contents were vary which depends on the individual and age of the plant. Diosgenin content of yam family were vary depending on the species and extraction methode.

According Behera et al.²², edible yams did not contribute for steroid production and the present study revealed that diosgenin content were very low in right twining species as compared to left twining species. The highest amount of diosgenin was observed in *D. bulbifera* (13.83 g/kg) followed by *D. hispida* (8.25 g/kg) and *D. pentaphylla* (8.18 g/kg). Among right twining species *D. oppositifolia* contained the highest amount of

diosgenin 6.58 g/ kg and among others *D.alata* had the lowest amount of diosgenin 0.95 g/ kg ascended by *D.belophylla* (1.20 g/ kg), *D.wallichii* (1.29 g/ kg).

Figure 2: Spectrogram of water soluble polysaccharides (WSP) from purple (a) and yellow (b) water yams



Water Soluble Polysaccharides (WSP)

Figure 2 showed a similar pattern of WSP purple and yellow water yams based on FTIR spectrum. It means that purple and yellow water yams had almost identical functional groups. Purple water yams were marked C-O-C functional groups at 1070.42 and 1105.14 cm⁻¹ while the yellow was at 1066.56, 1105.13, and 1145.65 cm⁻¹. C-O-C functional groups showed that a water soluble polysaccharides in purple and yellow water yams were -1, 4 mannosidic that connecting mannose foreach other. Functional group C=O stretch was shown in wave numbers 1652 and 1739.67 cm⁻¹. C=O group showed the aldehyde group making up a simple sugar mannose. It was supported Yu *et al.*²³ that the existence of a bond -1, 4 glucosidic and -1,4 mannosidic of glucomannan were marked with C-O-C stretch vibration at 1027.02 and 1244 cm⁻¹.


Results of this analysis indicated the presence of an acetyl group (CH₃-CO), respectively the number frequency 1242.07 cm⁻¹ and 1244 cm⁻¹ of WSP purple and yellow water yam, which was the side groups of glucomannan, hence it were suggested that the water soluble polysaccharide of water yam were glucomannan. According to Parry²⁴, glucomannan had an acetyl group by 10-19 units of carbon clusters at C2, C3, and C6 positions. Numbers wave frequency 2875.67; 2935.46, and 2962.46 cm⁻¹ in purple water yam whereas 2877.59; 2935.45, and 2962.46 cm⁻¹ in yellow water yam were a carboxylic acid functional group (RCOOH). Presence of carboxylic acid showed a WSP had the result of oxidation glucose or galactose. Simple sugars had primary alcohol group (carbon atom, C number 6) were changed to carboxyl, such as gluconic acid and galacturonic.

The wave number of 3406.06 cm⁻¹ in purple water yam and 3407.98 cm⁻¹ in yellow water yam were functional group of N-H (secondary amide, CONHR), that indicated the WSP was bound to proteins. This finding was supported by the result of previous research⁹ that dioscorin (a type of protein in *Dioscorea*) was strongly bound to the WSP from *Dioscorea hispida*. Dennst and Lu²⁵ revealed that dioscorin protein which was bound to the WSP had secondary structure.


The analysis using HPLC (KNAUER) and Aminex HPX 87H column. The column could be elute with standard sugars glucose, mannose, galactose, arabinose, rhamnose, glucuronic acid, and galacturonic acid. But, the type of sugar glucose and mannose could not be separate well as retention time almost the same. Figure 3, 4, and Table 3 showed the sugar analysis of WSP hydrolyzate from purple and yellow water yam. Hydrolyzate were obtained from dried WSP of purple or yellow water yam as much as 0.1 grams then were hydrolyzed with H_2SO_4 80% and 20%, neutralized with $Ba(OH)_2$ to pH 7 with final volume of 175 mL. WSP hydrolyzate contained more glucose. Mannose, arabinose, glucuronic acid and galacturonic acid in small quantities. Galactose and rhamnose were not detected in the hydrolyzate. Sugar analysis were affected polysaccharide hydrolysis process and the purity of WSP. $Ba(OH)_2$ could be reduce mannose level. Aman and Westerlund²⁶ explained the glucomannan and mannan could be precipitate by $Ba(OH)_2$ group likely due to the reaction of 2,3-cis-hydroxyl on mannose residues.

Dioscorea opposita contains mucus from mannan and proteins³. It is also supported by Liu *et al.*²⁷ that the major sugar composition of *Dioscorea batatas* is mannose, in addition to a small proportion of glucose and galactose in a ratio of 76.7, 16.9, and 6.3%, respectively. The sugar linkage of the polysaccharides was mainly 1,4-mannosidic. In addition, the acetyl groups are found in the polysaccharides, and the ratio of *O*-acetylation is estimated as 28% and shows the *O*-acetylation of C-2 and C-3 positions.

Figure 3: Chromatogram of glucuronic acid, galacturonic acid, glucose, arabinose (a) and mannose (b) from hydrolyzate WSP of purple water yams by AMINEX HPX 87H Column



a



b

Table 3: The results of sugar analysis from WSP hydrolyzate of purple and yellow water yam

Types of sugar	Purplewater yam		Yellowwater yam	
	Concentration (%)	% Relative	Concentration (%)	% Relative
Glucuronic Acid	0	0.00	0.002	5.71
Galacturonic Acid	0.001	2.56	0.001	2.86
Glucose	0.031	79.49	0.027	77.14
Galactose	0	0.00	0	0.00
Mannose	0.004	10.26	0.003	8.57
Rhamnose	0	0.00	0	0.00
Arabinose	0.003	7.69	0.002	5.71
Totale	0.039	100.00	0.035	100.00

* Concentration of glucuronicacidinpurple water yamwere0.0003% but rounded to0%

Figure 4: Chromatogram of glucuronic acid, galacturonic acid, glucose, arabinose (a) and mannose (b) from hydrolyzate WSP of yellow water yams by AMINEX HPX 87H Column





Figure 5: SDS PAGE of water yam (M:marker; A: purple water yam; and B: yellow water yam).




Figure 6: Chromatogram of amino acid purple water yam





Figure 7: Chromatogram of amino acid yellow water yam

Dioscorin

The water soluble protein in purple and yellow water yams contained dioscorin with molecular weight of 28.51 kDa. This finding was in accordance to the result that reported by Myoda *et al.*³ who indicated that molecular weight of *Dioscorea opposita* dioscorin were 23 and 32 kDa, while Gaidamashvili²⁸ revealed that molecular weight of dioscorin of *Dioscorea batatas* was 31 kDa.

The dioscorin content of purple and yellow water yams were 28.94% and 25.45% from total water soluble protein. The other types of proteins such as maltose binding lectin (68.68 kDa and 66.76 kDa), alcohol dehydrogenase (53.21 kDa), arabinogalaktan (16.17 kDa), and mannosebinding lectin (8.19 kDa) were found in *Dioscorea alata*. Gaidamashvili²⁷ found four major proteins of *Dioscorea batatas*, namely mannosebinding lectin (10 kDa), maltose binding lectin (66 kDa), dioscorin (31 kDa) and acidic chitinase homologous to chitinase from *Dioscorea japonica* with a ratio of 20:50:20:10. Protein band with a molecular weight of 16.7 kDa were arabinogalactan protein²⁹.

The protein of yellow and purple water yams were composed of aspartate, glutamate, serine, histidine, glycine, arginine, alanine, tyrosine, valine, phenylalanine, ileusin, leucine, and lysine by analysis HPLC and acid hydrolysis methode. Myoda *et al.*³ showed that dioscorin sequence consists of VEDEFSYIEGNPNPENWGN. The results of chromatogram showed that the peak of amino acids aspartate, glutamate, serine, glycine, tyrosine, valine, phenylalanine, and ileusin, supports the existence of protein dioscorin on yellow and purple water yams. However asparagine, tryptophan, and proline which were constituent of dioscorin, but it were not detected. Acid hydrolysis in sample preparation caused the damage of several other amino acids.

Conclusion

Purple water yam (*Dioscorea alata* var *Purpurea* and yellow water yam (*Dioscorea alata* L.) had different physical and chemical composition. The two tubers also contain bioactive compounds of diosgenin, water soluble polysaccharides, and dioscorin. Water soluble polysaccharides and soluble proteins from purple and yellow water yam had similar characteristics, although their levels were slightly different. Water soluble polysaccharide had methyl group (-CH₃), methylene (-CH₂), OH, NH, C=O, acetyl (C-O), carboxyl (COOH), and C-O-C. WSP hydrolyzate contained more glucose. Mannose, arabinose, glucuronic acid and galacturonic

acid in small quantities. Soluble protein of purple and yellow water yam contains dioscorin as one of bioactive compounds of yam family.

Acknowledgment

The authors would like to thank to Directorate General for Higher Education, Ministry of Education and Culture, Republic of Indonesia and Brawijaya University for supporting this research through Postgraduate Research Grant, DIPA of Brawijaya University No. 0636/023-04.2.16/15/2012.

References

1. Sulistyono E. and Jaminton M., Characterization of Tubers and Nutrition Contents from *Dioscorea* spp., Agronomy Bulletin, 2004, 32(2), 39-43.
2. Shah H.J. and Lele S.S. Extraction of Diosgenin, a Bioactive Compound from Natural Source *Dioscorea alata* var *Purpurea*, Journal Analytical and Bioanalytical Techniques, 2012, 3(4), 1-3.
3. Myoda T., Matsuda Y., Suzuki T., Nakagawa T., Nagai T., and Nagashima T., Identification of Soluble Proteins and Interaction with Mannanin Mucilage of *Dioscorea opposita* Thunb. (Chinese Yam Tuber), Food Sci. Technol. Res., 2006, 12(4), 299-302.
4. Kaimar, A., and Kathi, J.K. Wild Yam (*Dioscoreaceae*), The Longwood Herbal Task Force., The Center for Holistic Pediatric Education and Research., <http://www.mcp.edu/herbal/default.htm>, 1999.
5. Corbiere C., Liagre B., and Bianchi A., Different Contribution of Apoptosis to the Antiproliferative Effects of Diosgenin and other Plant Steroids, Hecogenin and Tigogenin, on Human 1547 Osteosarcoma Cells, Int J Oncol., 2003, 22, 899-905.
6. Trowel H., Definition of Dietary Fiber and Hypothesis That It Is a Protective Factor for Certain Diseases. Am J Clin Nutr., 1976, 29, 417-427.
7. Ha M.A., Jarvis M.C. and Man J.L., A definition for Dietary Fiber. Eur J Clin Nutr., 2000, 54, 861-864.
8. Chen H.L., C.H. Wang, C.T. Chang, and T.C. Wang, Effect of Taiwanese Yam (*Dioscorea japonica* Thunb. var *pseudojaponica* Yamamoto). on upper Gut Function and Lipid Metabolism in Balbic Mice, Basic Nutritional Investigation Nutrition, 2003, 19, 646-651.
9. Rachman F., Antihypertensive Effects of Water Soluble Polysaccharide Binding Dioscorin from Gadung (*Dioscorea hispida* Dennst.), Bachelor Thesis, Food Science and Technology, Agriculture Technology Faculty, Brawijaya University, Malang, Indonesia, 2011, 1-3.
10. Hou W.J., Liu J.S., Chen H.J., Chen T.E., Chang C.F., and Lin Y.H., Dioscorin, the major tuber storage protein of yam (*Dioscorea batatas* Decne) with carbonic anhydrase and trypsin inhibitor activities, J. Agric. Food Chem., 1999, 47, 2168-2172.
11. Hou W.C., Lee M.H., Chen H.J., Liang W.L., Han C.H., Liu Y.W. and Lin Y.H., Antioxidant Activities of Dioscorin, The Storage Protein of Yam (*Dioscorea batatas* Decne.) Tuber, J. Agric. Food Chem., 2001, 49, 4956- 4960.
12. Liu Y.W., Shang H.F., Wang C.K., Hsu F.L., and Hou W.C., Immunomodulatory Activity of Dioscorin, The Storage Protein of Yam (*Dioscorea alata* cv. Tainong No. 1) Tuber. Food Chem. Toxicol., 2007, 45, 2312- 2318.
13. Trivedi P.D, Kilambi P., Shivprakash R., and Karishma S.S., A Validated Quantitative Thin Layer Chromatographic Method for Estimation of Diosgenin in Various Plant Samples, Extract, and Market Formulation, Journal of AOAC International, 2007, 90(2), 358-363.
14. Herlina, Characterization, Hypolipidemic and Prebiotic Activity of Water Soluble Polysaccharide from Lesser Yam (*Dioscorea esculenta*), Dissertation, Postgraduate Program, Brawijaya University, Malang, Indonesia, 2012, 164-174.
15. Association of Official Analytical Chemists (AOAC), Association of Official Methods of Analysis (18 Edn). Official Analytical Chemist Inc. Mayland. USA, 2005.
16. Gracula M.C., Descriptive Sensory Analysis in Practice, Food and Nutrition Press Inc. Trumbull. Connecticut, 1997, 77, 5924-5931.

17. AspN.S, Claes G.J., Haakan H., and Monica S.,Rapid Enzymic Assay of Insoluble and Soluble Dietary Fiber,J. Agric. Food Chem.,1983, 31 (3), 476–482.
18. Liyana,PathirananC.M., dan ShahidiF., Antioxidan,Journal of BogorAgriculturalInstitute,2005,55-56.
19. OncinaR., Botía J., Del Río J. and Ortuño A., Analysis of Diosgenin. Food Chem.,2000,70, 489.
20. Charles A.L., HuangT.C., and Chang Y.H., Structural Analysis and Characterization of a Mucopoly saccharideIsolatedfrom Root of Cassava (*Manihot esculenaCrantz L*),Food Hydrocolloid,2008, 22, 184-191.
21. Lingga P., SarwonoB., RahardiF.,RahardjaP.C., AfriastiniJ.J., Wudianto R. and ApriadjiW.H., 1993, Planting Tubers., Penebar Swadaya. Jakarta, 52-53.
22. Behera K.K., SantilataS., andAratibalaP., Biochemical Quantification of Diosgenin and Ascorbic Acid from the Tubers of Different *Dioscorea* Species Found in Orissa,Libyan Agriculture Research Center Journal Internation, 2010, 1(2), 123-127.
23. YuH., Huang Y., Ying H., and XiaoC., Preparation and Characterization of a Quaternary Ammonium Derivative of Konjac Glucomannan, Carbohydrate Polym., 2007,69, 29-40.
24. Parry J.M., Konjac Glucomannan, In:Alan Imeson (ed). Food Stabilisers, Thickeners and Gelling Agents, John Willey and Sons. United Kingdom.,201, 198 – 216.
25. Lu Y.L., Chi C.Y., Liu Y.W., and HouW.C., Biological Activities and Applications of Dioscorins, the Major Tuber Storage Proteins of Yam,Journal of Traditional and Complementary Medicine,2011, 2 (1), 41-46.
26. AmanP. and Westerlund E., Cell Wall Polysaccharides Struktural, Chemical and Analytical Aspects in Carbohydrate in Food. Marcel dekker, Inc. Newyork,1996, 196.
27. Liu J.Y., Yang F.L., LuC.P., Yang Y.L., Wen C.L., Hua K.F., and WuS.H.,Polysaccharides from *Dioscorea batatas* Induce Tumor Necrosis Factor- α Secretion via Toll-like Receptor 4-Mediated Protein Kinase Signaling Pathways. J. Agric. Food Chem.,2008, 56,9892–9898.
28. Gaidamashvili M., Ohizumi Y., Iijima S., TakayamaT., OgawaT.,and Muramoto K.,Characterization of the Yam Tuber Storage Proteins from *Dioscoreabatatas* Exhibiting Unique Lectin Activities, The Journal of Biological Chemistry,2004,279(25), 26028-26035.
29. Du H., Simpson R.J., Clarke A.E., and Basic A., Molecular Characterization of a Stigma Specific Gene Encoding and Arabinogalactan Protein (AGP) from *Nicotiana glauca*, The Plant Journal,1996, 9, 313-323.

★★★★★