

Efficacy of baker's yeast as immunostimulant in nile tilapia (*Oreochromis niloticus*)

Henky Manoppo^{1*}, Usy N. Manurung², Reiny A. Tumbol¹

¹Laboratory of Fish Health, Environment and Toxicology, Faculty of Fisheries and Marine Science Sam Ratulangi University, Manado-Indonesia

²Department of Aquatic Science, Faculty of Fisheries and Marine Science Sam Ratulangi University Manado – Indonesia

Abstract: This study was conducted to evaluate the effect of baker's yeast cell as immunostimulant on nile tilapia (*Oreochromis niloticus*) immune responses. Fish was obtained from Aquaculture Development and Training Board (BP3I) Tateli, Fisheries and Marine Office of North Sulawesi and distributed into five outdoor concrete tanks at a density of 45 fish. After acclimatization for one week, fish was graded (mean weight 28.78 g) and set at 30 fish per tank. Fish in tank A received pellet not supplemented with baker's yeast while fish in tanks B, C, D, and E received pellet supplemented with 5, 10, 15, 20 g yeast cells/kg feed respectively. Feeding was applied four consecutive weeks at 5% of body weight per day, twice daily at 08.00 am and 16.00 pm. Immune parameters including total leucocytes count and phagocytosis activity were measured at week two and week four. After feeding for four weeks, administration of yeast cell exhibited significant effect on the total leucocyte count and phagocytosis activity ($p < 0.01$). The highest value of these two parameters was observed in fish treated with 5 g yeast cells. The result also showed that at high dose, the immune parameters declined as the administration time elongated. In conclusion, incorporation of baker's yeast cells into feed improved immunity of nile tilapia cultured outdoor by increasing total leucocyte count and phagocytosis activity of phagocyte cells.

Keywords: *Saccharomyces cerevisiae*, immunostimulant, nonspecific immune response, total leucocyte count, phagocytosis activity, *Oreochromis niloticus*.

Introduction

Nile tilapia (*Oreochromis niloticus*) is one of the important freshwater aquaculture species in Indonesia. This species grows relative fast and has high tolerance to environmental changes. Rapid intensification of aquaculture has raised the occurrence of infectious diseases weather caused by virus, bacteria, fungi and parasites. Disease problem has led to significant economic losses, continued to occur and probably has become the most limiting factor in tilapia aquaculture.

Improving fish performance and disease resistance of culture species to various pathogens are major challenges faced by fish culturist¹. Several attempts have been implemented to control the disease including the use of antibiotic and vaccine. However, the intensive use of chemicals or antibiotics in intensive aquaculture has brought about unwanted developments such as bioaccumulation, pollution, antibiotic-resistant pathogens, immunosuppression, and high expenditure^{2,3}. Antibiotic residue could be accumulated in the fish body and dangerous for human health. Antibiotic resistance can be transferred to the aquaculture environment and to human pathogens⁴. Vaccine has been proved to be effective in preventing the occurrence of diseases in aquaculture. However, the efficacy of vaccines was limited only on specific pathogen.

The use of immunostimulants offer an alternative to antibiotic or chemicals currently being used to combat disease in aquaculture has attracted more attention from researchers⁵. In contrast to vaccines, immunostimulant works by enhancing the non-specific immune response of fish and crustacean^{6,7}. Immunostimulant did not leave any residue in fish body and environment and not harmful for human health.

More environment-friendly disease control strategies are urgently needed to promote sustainable aquaculture production. Currently numerous natural products have been used as immunostimulant sources to control fish diseases such as herbals⁸⁻¹³, yeast^{3,14-17}, seaweed¹⁸. Herbs are currently used in commercial aquaculture as growth-promoting substances, antimicrobial agents, nutrients as well as many other applications. Their potential to prevent and control fish diseases is also being studied. Modulation of the immune response using medicinal plant products as a possible therapeutic measure has become the focus of extensive scientific investigation⁵.

Yeast products are frequently used as feed ingredients in aquaculture because of their nutritional value, which include proteins, lipids, vitamins and minerals. The major component of yeast cell wall is β -1-3 glucan (50–60%) capable of stimulating the non-specific immune function of fish and crustaceans^{3,17}. This research was carried out in October – December 2014 aiming to evaluate the effect of baker's yeast as immunostimulant on Nile tilapia immune responses.

Experimental

Fish used

Juveniles of Nile tilapia with an average body weight of 28.78 g were derived from Aquaculture Development and Training Board (BP31) Tateli, Fisheries and Marine Office of North Sulawesi. Fish were put in oxygenated-plastic bag, transported to the Faculty of Fisheries and Marine Science and then distributed into five 2x1x1 m³ outdoor concrete tanks (tank A, B, C, D and E) at a density of 45 individuals per tank. Each tank was equipped with one inlet pipe, outlet, and aerator. Fish were acclimatized for one week and fed standard pellet twice a day at 08.00 am and 16.00 pm at a rate of 5% of body weight per day.

Yeast

The research used baker's yeast (*Saccharomyces cerevisiae*) as immunostimulant bought from department store while feed used was commercial fish pellet.

Feed preparation

Standard pellet was supplemented with baker's yeast cells as immunostimulant. The doses of yeast were 5, 10, 15 and 20 g/kg of pellet while the control food was not supplemented with yeast cells. After being weighed, yeast cells were suspended in some pure water (1:10 v/w) and mixed thoroughly into pellet and air-dried at room temperature. After dry, feed was placed in plastic bag and stored at refrigerator at 4°C until use.

Research procedure and data collection

After acclimatization, fish was graded and set at 30 individuals per tank. Fish in tank A received pellet not supplemented with baker's yeast while fish in tanks B, C, D, and E received pellet supplemented with 5, 10, 15, 20 g yeast cells/kg feed respectively. Feeding was applied four consecutive weeks at 5% of body weight per day, twice daily at 08.00 am and 16.00 pm. During the experiment, water quality was kept stable by regular monitoring. To maintain at optimal level of water, water exchange as much as one-third was conducted once every three days.

Immune parameters including total leucocytes count and phagocytosis activity were measured at two weeks interval. Fish blood was withdrawn from vena caudalis according to the procedure of Stolen¹⁹. Blood sample was collected from three fishes of each treatment using a 1 mL syringe previously rinsed with anti-coagulant (EDTA) and transferred to sterile eppendorf. Measurement of immune parameters were conducted at day 14 and 28.

Total Leucocyte Count

The fresh blood samples were used for the calculation of total leucocytes and phagocytosis activity of phagocytes cells. To estimate the total number of leucocytes, 50 μ L of blood was placed in an appendorf and then mixed with 450 μ L of Turk's solution (1:10). The mixture was homogenized by gentle swing and incubated for five minutes at room temperature. Total leucocyte count was calculated using haemocytometer under light microscope.

Phagocytosis activity was estimated by mixing 0.1 mL of blood with an equal volume of zymosan as stimulator. The mixture of blood and yeast cells was homogenized by gentle movement and then incubated at room temperature for 20 minutes. Furthermore, the blood was smeared on glass slide and stained with Giemsa. Phagocytosis activity (%) was expressed as:

$$\frac{\text{Number of leucocytes engulfing zymosan}}{\text{number of leucocytes observed}} \times 100$$

Statistical analysis

Data obtained were expressed as mean \pm Stdv and analyzed by one-way analysis of variance (ANOVA). The difference effect between means was determined and compared by Duncan Test. Significant level was set at 0.05

Result and Discussion

Total Leukocyte

Oral administration of baker's yeast for two weeks exhibited non-significant effect on the total leucocyte count ($p=0.25$). Total leucocytes of fish between different treatment were almost similar. However, administration of baker's yeast showed significant effect ($p<0.01$) after being fed for four weeks (Table 1). The highest total leucocyte was achieved on fish fed feed supplemented with yeast cells at 5 g/kg feed, followed by 10 g/kg feed.

Table 1. Total leucocyte count of fish ($\times 10^7$ cells/mL) fed diets supplemented with different doses of yeast cells

Yeast cells (g/kg feed)	Week-2	Week-4
0	4.62 ^a	4.89 ^a
5	5.55 ^a	6.62 ^b
10	6.09 ^a	6.57 ^b
15	5.82 ^a	5.91 ^b
20	5.68 ^a	4.97 ^a

Different super scribes in the same column were significantly different

Phagocytosis Activity

Supplementation of yeast cells in feed for two weeks showed significant effect on phagocytosis activity of phagocytes cells of fish ($p=0.04$). This effect was sustained until weeks four of feeding ($p<0.01$). The highest phagocytosis activity of phagocytes cells was observed in fish fed feed with 5 g yeast cells per kg feed followed by 10 g/kg feed (Table 2).

Table 2. Mean phagocytosis activity (%) of nile tilapia fed on feed supplemented with baker's yeast cells

Yeast cells (g/kg feed)	Week- 2	Week- 4
0	27.97 ^a	32.37 ^a
5	35.98 ^{ab}	48.63 ^c
10	40.39 ^b	46.41 ^{bc}
15	41.08 ^b	42.66 ^b
20	45.01 ^b	34.72 ^a

Different super scribes in the same column were significantly different

Research results displayed that application of baker's yeast cells into feed could stimulate the increase of total leucocytes and phagocytosis activity of Nile tilapia phagocytes. It had been also reported that the use of yeast cell walls was able to enhance the innate immunity of Rohu (*Labeo rohita* Ham) and had a positive correlation with growth parameters²⁰. Through the absorption of yeast wall particle the immune function of fish was stimulated. In laboratory research, total leucocyte and phagocytosis activity of Nile tilapia (mean weight 10.57 g) increased significantly after feeding for four weeks with pellet supplemented with 10 g yeast cells per kg of pellet²¹. In common carp (*Cyprinus carpio* L), significantly increased phagocytic activity and superoxide anion production in kidney cells, and resistance to a bacterial pathogen, were observed in the yeast extract-treated fish compared to non-treated fish². In shrimp (*Marsupenaeus japonicus*), immune system of shrimp fed with yeast extract-supplemented diet increased significantly compared with shrimp fed control diet²². Supplementation of live bakers' yeast cells into fry Nile tilapia diet induced growth performance, feed utilization and immunity, and is promising as an alternative method to antibiotics for disease prevention in tilapia aquaculture¹. Yeast by-product from baker's yeast industry may be used as feed supplement and has positive effect on nonspecific immune response and growth of several fish species²³.

Baker's yeast is a particularly important natural bio-product since it contains immunostimulating compounds such as β -glucan, nucleotides, mannan, oligosaccharides and chitin^{2,14,24,25}. Such compounds have the capability to enhance immune responses of various fish species⁶. Immunostimulants may directly initiate activation of the innate immune defense mechanisms acting on receptors and triggering intracellular gene activation that may result in production of antimicrobial molecules²⁶. It leads to an increase in various components of immunity such as phagocytic activity, complement activity, lysozyme and disease resistance as well as serum Ig level¹².

Nucleotides play important role in essential physiological and biochemical functions including encoding and deciphering genetic information, mediating energy metabolism and cell signaling as well as serving as components of coenzymes, allosteric effectors and cellular agonists²⁵. Nucleotides are synthesized de novo in most of the tissues, but some immune and intestinal cells lack this process and thus depend on exogenous supply. Supplementation of nucleotides might improve cellular and humoral immune responses of various fish as well as shrimp. Oral administration of nucleotides to fish daily for 3 days resulted in enhanced responses of phagocytic and nitro blue tetrazolium (NBT) activities in kidney phagocytic cells²⁷. This activation of kidney cells was observed for at least 10 days post-treatment. The serum complement and lysozyme activities also increased in fish treated with nucleotides. Furthermore, the number of *A. hydrophila* in nucleotide-treated fish significantly decreased in the blood, kidney and liver after intraperitoneal injection. Thus yeast nucleotides appear to enhance nonspecific immune responses in fish. Nucleotides added to fish diet will optimize the function of cell replication including immune cells²⁸. Nucleotides increased immune responses and resistance of fish a number of pathogen simultaneously²⁹. Furthermore, supplementation of nucleotides also increased efficacy of vaccination, fish growth, larval quality and tolerance to stress. In grouper (*Epinephelus malabaricus*), fish fed diet supplemented with nucleotides for eight weeks had better growth and immune responses compared to control fish³⁰. In shrimp, supplementation of nucleotides in feed significantly enhanced immunity and resistance of *Litopenaeus vannamei*³¹.

β -glucan presents at the cell wall of yeast *S. cerevisiae* have a potent stimulatory effect on immune system of fish, crustacean and mammals^{6,32}. β -1,3/1,6-glucans bind specifically to a "receptor molecule" on the surface of phagocytes. After binding, the cells become more active in engulfing, killing and digesting bacteria and at the same time they secrete signal molecules (*cytokines*) which stimulate the formation of new white blood cells. And also activate antibody-producing white blood cells (B- and T-cells)^{7,33}. Supplementation of baker's yeast also increased feed and protein digestibility and thus resulted in better growth and feed efficiency³⁴.

Nile tilapia fed with pellet supplemented with β -glucan extracted from yeast *S. cerevisiae* significantly enhanced nonspecific immune response and resistance to *A. hydrophila*³⁵. In common carp (*Cyprinus carpio*), fish treated with β -glucan 0.1% had higher total leucocyte count and resistance to bacterial pathogen as compared to untreated fish³⁶. *Labeo rohita* (mean weight 35 ± 5 g) fed diet with β -glucan supplementation for 28 days had total leucocyte count higher than that of control fish³⁷. Another research found total leucocytes, lymphocytes and eosinophyl of Nile tilapia treated with *Saccharomyces* (10 g.kg⁻¹ feed), β -glucan (0.1%) and laminarian (0.1%) significantly decreased as the survival of fish decreases after exposed to mercury as stressor³⁸. Application of β -glucan by intraperitoneal injection on carp significantly enhanced immune responses and survival rate after challenged with *A. hydrophila*³⁹. Supplementation of β -glucan to the koi for 56 days showed considerable improvement in the immune response, growth, and survival of koi³².

In this research, there was no significant effect of yeast on total leucocytes and phagocytosis activity two weeks after feeding. As the doses of yeast increased the immune parameters increased too. But after feeding for four weeks, total leucocytes and phagocytosis activity of fish tended to decrease as the doses of yeast cells increased. At high doses, both total leucocytes and phagocytosis activity of fish treated with pellet contained 20 g of yeast cells/kg feed were low and almost similar with that of fish treated with control pellet. The best effect was observed in fish treated with low dose (5 g/kg of feed). This finding explained that the dose and administration time should be taken into account in applying an immunostimulant in aquaculture. The efficacy of immunostimulant by oral method decreases with long-term administration and overdoses of immunostimulants induce immunosuppression in fish⁶. Thus for the effective use of immunostimulants, dosages, method of administration, administration time and the physiological condition of fish need to be considered. In health management, dose and frequency of administration of immunostimulants are essential¹⁷.

Conclusion

Based on the results, it could be concluded that incorporation of baker's yeast cells into feed improved immunity of Nile tilapia cultured outdoor by increasing total leucocyte count and phagocytosis activity of phagocyte cells. The best effect was observed at low dose, but at high dose, the immune parameters declined as the administration time elongated. Thus research to establish the optimal administration time is necessary to be conducted.

Acknowledgements

The authors are grateful to the management of Aquaculture Development and Training Board (BP3I) Tateli, Fisheries and Marine Office of North Sulawesi for the supply of Nile tilapia juveniles.

References

1. Abdel-Tawwab M, Abdel-Rahman AM, Ismael NEM. Evaluation of commercial live bakers' yeast, *Saccharomyces cerevisiae* as a growth and immunity promoter for fry Nile tilapia *Oreochromis niloticus* (L) challenged *in situ* with *Aeromonas hydrophila*. Aquaculture, 2008, 280: 185-189.
2. Biswas G, Korenaga H, Takayama H, Kono T, Shimokawa H, Sakai M. Cytokine responses in the common carp, *Cyprinus carpio* L. treated with baker's yeast extract. Aquaculture, 2012, 356-357: 169-175.
3. Babu DT, Antony SP, Joseph SP, Bright AR, Philip R.. Marine yeast *Candida aquatextoris* S527 as a potential immunostimulant in black tiger shrimp *Penaeus monodon*. Journal of Invertebrate Pathology, 2013, 122: 243-252.
4. Wu YR, Gong QF, Fang H, Liang WW, Chen M, He RJ. Effect of *Sophora flavescens* on non-specific immune response of tilapia (GIFT *Oreochromis niloticus*) and disease resistance against *Streptococcus agalactiae*. Fish & Shellfish Immunology, 2013, 34 : 220- 227.
5. Galina J, Yin G, Ardo L, Jeney Z. The use of immunostimulating herbs in fish. An overview of research. Fish Physiol Biochem, 2009, 35: 669-676.
6. Sakai M. Current Research Status of Fish Immunostimulant. Aquaculture, 1999, 172 : 63-92.
7. Raa J. The use of immune-stimulants in fish and shellfish feeds. University of Tromsø Norway, 2000, 47-65.
8. Kumar S, Raman RP, Pandey PK, Mohanty S, Kumar A, Kumar K. Effect of orally administered azadirachtin on non-specific immune parameters of goldfish *Carassius auratus* (Linn. 1758) and resistance against *Aeromonas hydrophila*. Fish & Shellfish Immunology, 2013, 34 : 564-573.
9. Bilen S, Bulut M, Bilen A.M. Immunostimulant effects of *Cotinus coggyria* on rainbow trout (*Oncorhynchus mykiss*). Fish & Shellfish Immunology 2011, 30: 451-455.
10. Awad E, Austin B.. Use of *Lupinus perennis*, mango, *Mangifera indica*, and stinging nettle, *Urtica dioica*, as feed additive to prevent *Aeromonas hydrophila* infection in rainbow trout *Oncorhynchus mykiss* (Walbaum). Journal of Fish Diseases, 2010, 33: 413-420.
11. Punitha SMJ, Babu MM, Sivaram V, Shankar VS, Dhas SA, Mahesh TC, Immanuel G, Citarasu T.. Immunostimulating influence of herbal biomedicines on nonspecific immunity in Grouper *Epinephelus tauvina* juvenile against *Vibrio harveyi* infection. Aquacult Int, 2008, 16: 511-523.

12. Divyagnaseswari M, Christyapita D, Michael RD. Enhancement of nonspecific immunity and disease resistance in *Oreochromis mossambicus* by *Solanum trilobatum* leaf fractions. *Fish & Shellfish Immunology*,2007, 23: 249-259
13. Yin G, Jeney G, Racs T, Xu P, Jun X, Jeney Z. Effect of two Chinese herbs (*Astragalus radix* and *Scutellaria radix*) on nonspecific immunes system of tilapia, *Oreochromis niloticus*. *Aquaculture*,2006, 253:39-47.
14. Reyes-Becerril M, Tovar-Ramirez D, Ascensio-Valle F, Civera-Cerecedo R, Gracia-Lopez V, Barbosa-Solomieu V. Effects of dietary live yeast *Debaryomyces* on the immune and antioxidant system in juvenile leopard grouper *Mycteroperca rosacea* exposed to stress. *Aquaculture*,2008, 280: 39-44
15. Sarlin PJ, Philip R. Efficacy of marine yeast and baker's yeast as immunostimulants in *Fenneropenaeus indicus*: A comparative study. *Aquaculture*,2011, 321: 173-178
16. Ma Y, Liu Z, Yang Z, Li M, Liu J, Song J. Effects of dietary life yeast *Hanseniaspora opuntiae* C21 on the immune and disease resistance against *Vibrio splendidus* infection in juvenile sea cucumber *Apostichopus japonicus*. *Fish & Shellfish Immunology*, 2013, 34: 66-73
17. Sajeevan TP, Philip R, Singh ISB. Dose/frequency: a critical factor in the administration of glucan as immunostimulant to Indian white shrimp *Fennerppenaeus indicus*. *Aquaculture*,2009, 287: 248-252
18. Ganeshamurthy R, Dhayanithi NB, Kumar TTA, Kumaresan S. Evaluation of antibacterial activity and immunostimulant of red seaweed *Chondrococcus hornemanni* (Kuetzing, 1847) against marine ornamental fish pathogen. *Journal of Coastal Life Medicine*, 2014, 2(1): 64-69
19. Stolen SJ. Techniques in immunology. 1st edition. SOS Publication 43 de Normandie A Venue Fair Haven, NJ, 1990.
20. Tewary A, Patra BC. Oral administration of baker's yeast (*Saccharomyces cerevisiae*) acts as a growth promoter and immunomodulator in *Labeo rohita* (Ham.). *Journal of Aquaculture Research and Development*, 2011, 2:1-7
21. Manurung UN, Manoppo H, Tumbol RA.Evaluasi ragi roti (*Saccharomyces Cereviciae*) sebagai imunostimulan dalam meningkatkan respon imun non spesifik dan pertumbuhan ikan nila (*Oreochromis Niloticus*). *Jurnal Budidaya Perairan*, 2013, 1: 8-14
22. Biswas G, KorenagaH, Nagamine R, Kono T, ShimokawaH, Itami T, Sakai M.Immune stimulant effects of a nucleotide-rich baker's yeast extract in the kuruma shrimp, *Marsupenaeus Japonicus*. *Aquaculture*, 2012,366-367: 40-45.
23. Olivia-Teles A, Goncalves P. Partial replacement of fishmeal by brewer's yeast *Saccharomyces cerevisiae*, in diets for sea bass *Dicentrarchus labrax* juveniles. *Aquaculture*,2001, 202: 269'278
24. Li P, Galtin III DM. Evaluation of brewers' yeast. (*Saccharomyces cereviciae*) as a feed supplement for hybrid striped bass (*Marone chrysops x M. saxatillis*). *Aquaculture*, 2003, 219: 681-692
25. Li P, Galtin III DM. Nucleotide nutrition in fish: Current knowledge and future application. *Aquaculture*,2006, 251 : 141 – 152.
26. Bricknell I, DalmoRA. The use of immunostimulants in fish larval aquaculture. *Fish & Shellfish Immunology*,2005, 19: 457-472.
27. Sakai M, Taniguchi K, Mamoto K, Ogawa H, Tabata M. Immunostimulant effects of nucleotide isolated from yeast RNA on crap, *Cyprinus carpio* L. *J Fish Dis*, 2001, 24: 433-438
28. Sajeevan TP, Philip R, Singh ISB. Immunostimulatory effect of a marine yeast *Candida sake S156* on *Fenneropenaeus indicus*. *Aquaculture*,2006,257: 150-155.
29. Burrels C, Williams PD, FomoPF. Dietary nucleotide: A novel supplement in fish feed effects on resistance to disease in salmonids. *Aquaculture*,2001, 199 : 159 – 169
30. Lin YH, Wang H, Shiao SY. Dietary nucleotide supplementation enhance growth and immune response of grouper, *Epinephelus malabaricus*. *Aquaculture Nututrition*, 2009, 15: 117-122
31. Manoppo H, Sukenda, Djokosetyanto D, Fatuchri M, Harris E. Nukleotida meningkatkan respon imun dan performa pertumbuhan udang vaname, *Litopenaeus vannamei*. *Jurnal Aquacultura Indonesiana*, 2009,Vol. 10 (2): 85-92
32. Lin S, Pan Y, Luo L, Luo L. Effects of dietary β -glucan, chitosan or raffinose on the growth, innate immunity and resistance of koi (*Cyprinus carpio koi*). *Fish & Shellfish Immunology*,2011, 31: 788-794
33. Gannam AL, Schrock RM. Immunostimulant in fish diet *in* Nutrion and Fish Health. Food Products Press, New York, 2001, 235-260
34. Wache' Y, Auffray F, Gatesoupe FL, Zurrbonino J, Gayet V, Labbe' L, Quentel C. Cross effect of the strain dietary *Saccharomyces cereviciae* and rearing condition on the onset of intestinal micro biota and digestive enzymes in rainbow trout, *Onchorhynchus mykiss* fry. *Aquaculture*,2006, 258:470-478

35. Jamal IN. Penggunaan β -Glukan yang di Ekstrak Dari Ragi Roti *saccharomyces cerevisiae* Untuk Meningkatkan Sistem Imun Non Spesifik Ikan Nila (*Oreochromis Niloticus*). (Thesis). Pascasarjana Universitas Sam Ratulangi Manado-Indonesia, 2013.
36. Sahan A, Duman S. Effect of β Glucan on hematology of common carp (*Cyprinus carpio*) infected by ectoparasites. *Mediterranean Aquaculture*, 2010, (1); 1-7.
37. Misra CK, Das BK, Mukherjee SC, Pattnaik P. Effect of multiple injections of β -glucan on non-specific immune response and disease resistance in *Labeo rohita* fingerlings. *Fish & Shellfish Immunology*, 2006, 20: 305-319.
38. El-Boshy ME. Immunomodulatory effect of dietary *Saccharomyces cerevisiae*, β -glucan and laminaran in merciric chloride treated nile tilapia *Oreochromis niloticus* and experimentally infected with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 2010, 28 : 802 - 808.
39. Selvaraj V. Administration of yeast glucan enhances survival and some non-specific and specific immune parameters in carp *Cyprinus carpio* infected with *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 2005, 19 : 293 - 306.
