



Evaluation of water quality of the surface water of the Damietta Nile Branch, Damietta Governorate, Egypt

M.S.M. EL-Bady and H.I. Metwally

Department of Geological Sciences, National Research Center, Dokki, Cairo, Egypt

Abstract : Water quality and heavy metals pollution of the Damietta Nile Branch at Damietta governorate were investigated in the current work. fourteen different sampling points were selected along the Damietta Nile branch from Ras EL-Bar (sample 1) to Sheremsah (sample 14). Physical and chemical parameters and the concentrations of Cd, Cr, Cu, Ni, Fe, Al, Hg, Pb and Zn were investigated for water quality assessment of Damietta Nile Branch at Damietta Governorate. Most of water suitable for irrigation, while some of water suitable for drinking. The water near the Damietta Nile Branch Outlet is unsuitable for drinking and irrigation, but the water in the south of study area suitable for all purposes.

Keywords: Water Quality Indices – Damietta Governorate – Nile River – Pollution.

Introduction:

It is known that the 1959's Nile water agreement with Sudan, allocates 55.5 BCM (billion cubic meters)/ year to Egypt. Water of Nile River has been used in many of purposes such as drinking (domestic) water supply, irrigation, industrial, fisheries, recreation. Water requirements in Egypt are continuously increasing due to the high rates of growing population, increasing urbanization, industrialization and the rapid agricultural growth. Demands for the agricultural sector represent the largest component (about 80 %) of the total water demand in Egypt. According to ¹ the actual release from the High Aswan Dam shows very little yearly variation. Annual variation of the release of water from the High Aswan Dam depends mainly on irrigation needs. The release from the high Aswan Dam ranges from approximately 800 m³/s during the (winter) closure period to approximately 2760 m³/s during the summer months. In the Nile Valley and Nile Delta, groundwater resource use account for approximately 4.4 billion m³/yr., mainly being recharged from the Nile and from seepage from irrigated agriculture. Rainfalls plays a minor role in Egypt's water resources, with average rainfall rates declining from 200 million m³/yr. at the Mediterranean Coast to 20 mm³/yr. in Cairo and almost zero in Upper Egypt.

In the 1993/94 hydrological season, gross water consumption of irrigated agriculture amounted to approximately 54.5 B. (billion) m³/yr. (of which almost 30 B. m³/yr. occurred in the Delta). Other water users, such as municipal and domestic drinking water supply, industry and others consumed approximately 8.8 B. m³/yr., and an estimated 2.0 B. m³/yr. was lost through evaporation and about 14.0 B. m³/yr. was discharged to the Mediterranean Sea (Table 1). In recent years, the amount discharged to the Mediterranean Sea directly from the River Nile system, declined sharply, in favour of agricultural and drinking water supply use (approximately 2.3 B. m³ discharged in 1990 compared with approximately 1.2 B. m³) ¹.

Table (1): Water Budget Calculation of the River Nile System [1].

Resources	Billion m³/yr.	Water Diversions/ Intakes	Billion m³/yr.
Release From Lake Naser	55.470	Agricultural Intake	54.410
Groundwater Extraction	4.400	Drinking Water Supply	2.910
Irrigation Excess Water Return	16.910	Industrial Water Supply	5.890
Industrial Wastewater Return	5.480	Nile Flow to the Med. Sea	1.160
Domestic Wastewater Return	1.400	Drainage Outflow	12.890
		Evaporation From System	2.000
		Groundwater Recharge	4.400
Total	83.660	Total	83.660

Egypt suffering from increasing deterioration of its water due to increasing discharges of polluted domestic and industrial effluents into its waterways such as River Nile and Canals. The pollution of water due to the excessive use of pesticides and fertilizers in agriculture. Deterioration in water quality occurs when the Nile divided into Damietta and Rosetta Branches due to the disposal of the municipal and industrial effluents and agricultural drainage with decreasing flows².

Water quality is a term used to express the suitability of water to various purposes³. The quality of water may be described in terms of the concentration and state the organic and inorganic material present in the water, together with certain physical characteristics of the water⁴.

According to ¹ Egyptian industry uses 638 M. m³/yr. of water, of which 549 M. m³/yr. is discharged to the drainage system. Industrial activities in the Greater Cairo and Alexandria regions use 40% of the total. The River Nile supplies 65% of the industrial water needs and receives more than 57% of its effluents. More detailed information about water consumption, wastewater discharge and point sources of pollution and loads from different industrial sectors are provided (Table 2).

Table (2): Industrial Wastewater Discharge to the River Nile System District [1].

District	Ultimate Sink (M.m³/yr.)				Total
	Nile	Canals	Drains	Lakes	
Upper Egypt	192	5	2	5	204
Greater Cairo	80	21	20	7	128
Delta	27	85	13	1	126
Alexandria	13	7	33	35	88
Others	0	0	3	1	4
Total	312	118	71	49	550

The natural and anthropogenic effects have been changed the geo-environmental status of Nile Delta in the last few decades. The Nile Delta has ecological and economic values and it considered a major centre of population and agriculture⁵. Water quality has been changed by human intervention⁴. These effects are the polluting activities, such as the discharge of domestic, industrial, urban and other wastewaters into the watercourse (whether intentional or accidental) ⁶.

According to ^{7,8} 124 source points of pollution discharge the wastewater to the Nile River from Aswan to Delta Barrage, of which 67 are agricultural drains and the remainder is industrial and domestic sources. Industry purpose in Egypt uses 7.6 Bm³/yr of water. In the Delta region the amount of agricultural drainage water reuse officially was estimated to be around 4.27 BCM/year, in addition to about 0.3 BCM/year lifted to surface water (Rossetta branch) from west delta drains. Additional unofficial reuse done by farmers themselves has been estimated to be around 2.8 BCM/year. The remaining drainage water is discharged to the sea and the northern lakes via drainage pump stations. The total amount of drainage water that was pumped to the sea has been estimated to be 12.41 BCM/year⁸. The changes in water quality are primarily due to a combination of land and water use, as well as water management interventions such as; (a) different hydrodynamic regimes regulated by the Nile barrages, (b) agricultural return flows, and (c) domestic and industrial waste discharges including oil and wastes from passenger and riverboats⁷.

The situation of water pollution is probably getting worse with time, as the discharge of wastes is increasing. Heavy metals contamination in river is one of the major quality issues in many fast growing cities, because maintenance of water quality and sanitation infrastructure did not increase along with population and urbanization growth, especially for the developing countries^{9,10,11}. Heavy metals contamination is important due to their potential toxicity for the environment and human beings^{12,13,14,15}. Some of the metals such as Cu, Fe, Mn, Ni and Zn are essential as micronutrients for the life processes in animals and plants while many other metals such as Cd, Cr, Pb and Co have no known physiological activities^{16,17}. Damietta branch is about 242 km length, with average width and depth 200 and 12 meter respectively. It's the main source of drinking and irrigation waters for many of Governorates such as El-Qalubia, El-Gharbyia, El-Dakahlyia and Damietta¹⁸. Faraskour Dam divides the Damietta Nile branch in Damietta Governorate at the ilet of Damietta City about 20 km south of Mediterranean Sea to cut off the flow of the Nile water to the Mediterranean Sea. The water characteristics after the dam (saline water) is completely different compared with the water before the dam (fresh water)¹⁹. Damietta branch receives several pollutant types. Talkha fertilizer factory is considered the main source of industrial pollution, recooling waters of Talkha and Kafr Saad electric power stations. Domestic and sewage effluents at El-Serw City represent another source of pollution²⁰.

Damietta Nile Branch at Damietta Governorate has been suffered from intensive pollution. Damietta Nile Branch receives the water of a number of agricultural drains, which are heavily polluted by industrial and domestic sewage. The Damietta Branch receives polluted water of a number of agricultural drains, The Fertilizer Company is considered as the major point source of industrial pollution at Damietta branch. Many of villages on each banks of Damietta Nile branch without sanitation services, thus the river receive many wastewater resulted from industrial and domestic activities.

Damietta Governorate with area of about 910 km² north of the Nile Delta and contains about 1.3 million persons living within its four administrative centers. These centers are Kafr Saad, Faraskour, El-Zarka and Damietta center. It was the Egyptian gate along the Mediterranean Sea before the construction of Alexandria about 300 years BC. The long of the Damietta Nile Branch in the Damietta Governorate about 40 km from Ras EL-Bar (north) to Sheremsah Village (south). The landuse/cover map of Damietta Governorate was showed in figure 1. It consists of water, barren land, agriculture and urban²¹. About 14 water samples were taken from the Damietta Nile Branch from Ras EL-Bar to Sheremsah Village (Table 3) to study the water quality in the Nile Water of Damietta Governorate.




Fig.(1). The land use/cover map of Damietta Governorate as obtained from the Landsat-8 OLI image classification [21].

Materials and Methods:

Water samples were collected from the Damietta Nile branch at Damietta governorate from March to May 2015 and tested for physical qualities and chemical contents. The samples were collected from 14 sampling stations along the main flow of the Damietta Nile branch (Fig. 2, Table 3) starting from upstream near Sheremsah Village (sample 14) to downstream (outlet) at Ras El-Bar City (sample 1). Location of samples has been recorded using global positioning system (GPS). Water samples were collected from the central area of each site at depth of 10 – 30 cm.

Water samples were collected and kept into a one-litre polyethylene bottle in ice box and analyzed in the laboratory. Some of the physicochemical parameters including the electrical conductivity of the water samples ($\text{mS}\cdot\text{cm}^{-1}$), pH, and water temperature ($^{\circ}\text{C}$) were measured in the field by using water checker U-10 Horiba Ltd. The other water parameters were measured according to the traditional manual methods of the American Public Health Association^{22,23}. Total Pb, Ni, Al, Cu, Cr, Mn, Zn, Hg, Fe, Cd were measured after digestion using spectrophotometry.




Fig. (2). Location of the water samples of Damietta Nile Branch in Damietta Governorate

Table (3): Location of the samples along the Damietta Nile Branch course

Samples	Latitude	Longitude
1 (Ras ELBar)	31° 29' 33.80"	31° 49' 40.71"
2	31° 27' 52.18"	31° 48' 56.33"
3	31° 26' 2.18"	31° 48' 0.55"
4	31° 24' 37.68"	31° 47' 4.16"
5	31° 24' 41.01"	31° 45' 20.56"
6	31° 23' 7.59"	31° 44' 22.84"
7	31° 22' 52.64"	31° 42' 55.53"
8	31° 20' 40.22"	31° 42' 46.03"
9	31° 19' 13.18"	31° 41' 59.21"
10	31° 18' 9.68"	31° 41' 17.74"
11	31° 16' 57.63"	31° 41' 25.64"
12	31° 17' 25.89"	31° 39' 25.24"
13	31° 15' 12.99"	31° 40' 26.39"
14 (Sheremsah)	31° 14' 37.16"	31° 38' 48.64"

Water Quality Indices:

To investigate the water quality, four indices were used individually in this study, water quality index (WQI) using PH, TDS, Ec, DO, BOD, COD, CO₃, HCO₃, Cl, SO₄, Ca, Mg, NO₃, Hardness and alkalinity parameters, and Metal pollution index (MPI), Pollution Index (PI) and Heavy Metal Pollution Index (HMPI) using the heavy metals concentrations. Values of desirable and maximum allowable limits of different parameters, according to ^{24,25} for drinking water and ²⁶ for irrigation purpose are listed in Table 4. Each parameter is assigned a weight according to its relative importance for quality of water for drinking purposes, as shown in Table 1. Maximum weight of 5 is assigned to Total Dissolved Solids (TDS), PH, EC,(NO₃), (Mg), (Ca), (DO) hardness and alkalinity Pb, Cd, Hg, and weight of 4 is assigned to BOD, COD, Mn and Cr, weight of 3 is assigned to (SO₄), (Cl), and Fe, and weight of 2 is assigned (CO₃), (HCO₃), (NO₃), Al, Ni, Zn and Cu ²⁷ (Tables 5 and 6). The mathematical formula used to determined the water quality are as following:

Table (4): Standards of drinking and irrigation water according to WHO, FAO and Egyptian standards

Parameters	Mean	Drinking water standards			Irrigation standards
		[25]	Maximum	[24]	
Water					
PH	7.94	6.5-8.5		8.5	8.5
TDS mg/L	2565.48	1000	1000	500	2000
EC μ s/cm	4008.57	2000			3000
DO mg/L	7.87	6			
BOD mg/L	3.22	3			
COD mg/L	15.57	10		10	
CO ₃ ²⁻ mg/L	9.85	100		100	3
HCO ₃ ⁻	148.28	100		100	610
Cl- mg/L	1065.21	250	250	200	1063
SO ₄ ²⁻ mg/L	64.07	250	250	250	960
Ca ²⁺ mg/L	69	75	350	75	400
Mg ²⁺ mg/L	71	50	150	50	60
NO ₃ - mg/L	26.5	44	45	50	
T-	140	500	500	500	
Alkalinity	54.35			250	
Pb mg/L	0.020	0.01	0.01	0.01	5
Cd mg/L	0.002	0.003	0.003	0.003	0.01
Zn mg/L	0.191	3	3	0.5	2
Cumg/L	0.012	2	2	2	0.2
Cr mg/L	0.100	0.05	0.05	0.05	
Fe mg/L	0.221	0.3	0.3	0.3	5
Nimg/L	0.011	0.02	0.02	0.07	0.2
Mnmg/L	0.06	0.1	0.4	0.1	0.2
Al mg/L	0.017	0.2	0.2	0.2	5
Hg mg/L	0.001	0.001	0.001	0.001	

Table (5): Parameters of the water of Damietta Nile Branch in Damietta Governorate

Parameters	Samples														Relative Importance	Wi
	1	2	3	4	5	6	7	8	9	10	11	12	13	14		
PH	8.41	8.39	8.39	8.4	8.12	7.98	7.85	8.29	7.21	7.77	7.25	8.11	7.99	7.1	5	0.083
TDS mg/L	14369.28	9808.64	5483.52	3636.48	422.4	270.72	289.28	270.08	255.36	227.84	220.8	222.72	224	215.68	5	0.083
EC μ s/cm	22452	15326	8568	5682	660	423	452	422	399	356	345	348	350	337	5	0.083
DO mg/L	12.5	10.2	9.8	8.4	7.4	7.1	6.8	6.4	6.4	6.2	6.3	6.4	8.2	8.1	5	0.083
BOD mg/L	4.9	4.2	4.2	3.9	2.2	2.2	2.6	3.1	3.5	2.1	3.6	4.1	2.2	2.4	4	0.066
COD mg/L	28	24	25	16	12	14	15	11	13	12	12	14	11	11	4	0.066
CO ₃ ²⁻ mg/L	15	11	12	11	11	10	12	9	8	7	9	8	8	7	2	0.033
HCO ₃ ⁻ mg/L	221	198	200	172	156	125	124	127	122	142	122	121	124	122	2	0.033
Cl ⁻ mg/L	8653	5645	82	66	52	53	58	55	49	51	42	40	34	33	3	0.05
SO ₄ ²⁻ mg/L	290	123	101	66	52	41	32	28	24	33	25	28	25	29	3	0.05
Ca ²⁺ mg/L	322	240	101	29	33	23	34	22	23	28	28	29	31	23	5	0.083
Mg ²⁺ mg/L	421	320	15	20	19	22	25	24	24	23	22	20	21	18	5	0.083
NO ₃ ⁻ mg/L	17	18	15	19	20	21	20	19	29	32	36	41	40	44	2	0.033
T- Hardness mg/L	743	560	116	49	52	45	59	46	47	51	50	49	52	41	5	0.083
Alkalinity mg/L	126	121	102	85	30	35	42	33	36	38	31	25	29	28	5	0.083
Sum Wi																1

Table (6): Heavy metals concentration of the Damietta Nile Branch water in Damietta Governorate

Heavy Metals	Samples														Relative	Wi
	1	2	3	4	5	6	7	8	9	10	11	12	13	14		
Pb mg/L	0.03	0.04	0.03	0.04	0.02	0.02	0.01	0.01	0.001	0.01	0.01	0.04	0.01	0.01	5	0.147
Cd mg/L	0.003	0.003	0.004	0.004	0.002	0.001	0.003	0.002	0.001	0.001	0.001	0.001	0.002	0.002	5	0.147
Zn mg/L	0.4	0.3	0.2	0.5	0.01	0.2	0.06	0.005	0.02	0.4	0.06	0.5	0.01	0.01	2	0.058
Cu mg/L	0.02	0.02	0.02	0.03	0.01	0.01	0	0.01	0.01	0	0	0.03	0.01	0	2	0.088
Cr mg/L	0.04	0.05	0.04	0.9	0.02	0.03	0.05	0.06	0.05	0.04	0.04	0.06	0.02	0.01	4	0.117
Fe mg/L	0.4	0.2	0.3	0.5	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.5	0.1	0.1	3	0.058
Ni mg/L	0.01	0.01	0.02	0.02	0	0	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.01	2	0.058
Mn mg/L	0.09	0.08	0.06	0.1	0.02	0.08	0	0	0.08	0.06	0.04	0.05	0.09	0.09	4	0.117
Al mg/L	0.2	0.01	0.02	0.2	0.02	0.02	0.03	0.01	0.01	0.05	0.05	0.01	0.01	0.01	2	0.058
Hg mg/L	0.002	0.002	0.001	0.003	0.002	0.001	0.002	0	0.001	0.001	0.001	0.001	0	0	5	0.147
Sum Wi																1

(1) Water Quality Index (WQI):

WQI has been calculated to evaluate the suitability of water quality of Damietta Nile Branch using the weighted arithmetic water quality index method, which classifies the water quality according to the degree of purity by using the most commonly measured water quality variables. The calculation method of WQI was developed by²⁸, which has been widely used by many scientists^{29,30,31,32}. The grading of the water quality is showed in Table. 7.

$$WQI = \frac{\sum_{i=1}^n Q_i \times W_i}{\sum_{i=1}^n (W_i)}$$

Where Q_i (water quality rating) = $100 \times (V_a - V_i) / (V_s - V_i)$,

Where V_a = actual measured value of the water sample, V_i = ideal value (0 for all parameters except pH and DO which are 7.0 and 14.6 mg l-1 respectively). V_s = standard value.

$W_i = w_i / \sum w_i$ Where w_i is the weight of i th parameter and n is the number of chemical parameters.

Table.(7): Water Quality Index according to³³.

WQI Values	Grading
0–25	Excellent
26–50	Good
51–75	Poor
76–100	Very Poor
Above 100	Unsuitable for drinking purpose

(2) Metal Pollution Index:

Metal pollution index (MPI) is based on a total trend evaluation of the present status. It used to determine the metal contamination of Damietta Nile Branch Water. The higher the concentration of a metal compared to its respective MAC value, the worse the quality of the water. MI value >1 is a threshold of warning³⁴. According to³⁵ the MI is calculated by using the following formula:

$$MPI = \sum_{i=1}^n \left(\frac{C_i}{MAC_i} \right)$$

where: C_i : mean concentration

MAC: maximum allowable concentration

Water quality and its suitability for drinking purpose can be examined by determining its metal pollution index^{36,37}. According to³⁵ water samples can be divided into three groups including: potable ($MPI < 1$), on the threshold of danger of drinking ($MPI = 1$) and non-potable ($MPI > 1$).

The quality grading for irrigation water were modified according to heavy metals. Where, MPI divided into suitable for irrigation or no threshold of danger ($MPI < 1$), on the threshold of danger of irrigation ($MPI = 1$) and unsuitable for irrigation or high danger ($MPI > 1$).

(3) Pollution index (PI):

The pollution index was used in this study to evaluate the degree of heavy metal contamination in water samples^{38,39,40,41}. The tolerable level is the element concentration in the water considered safe for human consumption⁴². Pollution index (PI) is based on individual metal calculations and categorized into 6 classes (Table 8) according the following equation⁴³.

$$PI = \sum_{i=1}^n \frac{(C_i)}{S_i} \times Nm$$

Where C_i = Heavy metal concentration in water; S_i = permissible Level and Nm = Number of Heavy metals. Water sample with Pollution Index (PI) greater than 1 is regarded as being contaminated.

Table. (8): Categories of Water Pollution Index.

Class	PI value	Class
1	<1	No effect
2	1–2	Slightly affected
3	2–3	Moderately affected
4	3–5	Strongly affected
5	>5	Seriously affected

(4) Heavy Metal Pollution Index (HMPI):

The HMPI, represent the total quality of water with respect to heavy metals. HMPI was first suggested in 1996 that represents the overall quality of water which is based on heavy metals³⁶. The index is calculated based on the weighting the parameters that the weight value is between zero and one, points the importance of the parameters. This index is calculated by the following equation:

$$HMPI = \frac{\sum_{i=1}^n Q_i \times W_i}{\sum_{i=1}^n (W_i)}$$

Where Q_i (water quality rating) = $100 \times (V_a - V_i) / (V_s - V_i)$,

Where V_a = actual measured value of heavy metals of the water sample, V_i = ideal value of the heavy metals (0 for each heavy metals) V_s = standard value.

$W_i = w_i / \sum w_i$ Where w_i is the weight of i th parameter and n is the number of chemical parameters.

Water quality based on heavy metal pollution index can be divided into three categories including: low heavy metal pollution (HMPI <100), heavy metal pollution on the threshold risk (HMPI = 100) and high heavy metal pollution (HMPI > 100)³⁶. If the samples have heavy metal pollution index values greater than 100, water is not potable.

Results and Discussion**Physico-Chemical Characteristics :**

The physico-chemical parameters of Damietta Nile branch water in Damietta Governorate were shown in Tables 5 and 6 compared with the standards values of drinking purpose in Table 4. PH values range from 7.1 to 8.41 with mean value was 7.94. All values of PH were in the normal range of ^{24,25} (Table. 4). PH used as indicators of alkalinity and acidity of water, where, the PH values may affects in many chemical and biological processes in the water. TDS values range from 215.68 to 14369.28 mg/L with mean value was 2565.48 mg/L. TDS values were lower than the values of ^{24,25} in all samples except in samples 1,2,3 and 4. EC values ranged from 337 to 22452 μ S/cm, with mean of 4008.57 μ S/cm. EC values were lower than the values of ²⁵ in all samples except in samples 1,2,3 and 4. Dissolved Oxygen (DO) is required for the metabolism of aerobic organisms and it influences organic decomposition, where, DO was ranged from 6.2 to 12.5 mg/L with mean of 7.87 mg/L. DO values were more than the values of ²⁵ in all samples. The BOD ranged from 2.1 mg/L to 4.9 mg/L, with mean of 3.22 mg/L. Most of BOD of the samples more than ²⁵ except samples 5,6,7, 10, 13 and 14. The COD ranged from 11 mg/L to 28 mg/L, with mean 15.57 mg/L. All COD of samples more than the value of ²⁵. CO₃ concentration ranged between 7 mg/L to 15 mg/L with an average value of 9.85 mg/L while the concentration of HCO₃ varied from 121- 221 mg/L with a mean value of 148.22 mg/L. CL concentration ranged between 33 mg/L to 8653 mg/L with an average value of 1069.21 mg/L while the concentration of SO₄ varied from 24- 290 mg/L with a mean value of 64.07 mg/L. CO₃ concentration ranged between 22 mg/L to 322 mg/L with an average value of 69 mg/L, while the concentration of Mg varied from 18 - 421 mg/L with a mean value of 71 mg/L. NO₃ concentration ranged between 15 mg/L to 44 mg/L with an average value of 26.5 mg/L. Hardness ranged between 41 – 743 mg/L with average of 140 mg/L, while Alkalinity ranged from 25 to 126 mg/L, with mean of 54.4 mg/L. The concentration of these heavy metal (Pb, Cd, Zn, Cu, Cr, Fe, Ni, Mn, Al and Hg) were introduced into the Damietta Nile Branch in Damietta Governorate from agricultural drains, which are heavily polluted by industrial and domestic sewage, many of villages on each banks of Damietta Nile branch without sanitation services, Kafr Saad electric power stations, and Talkha fertilizer factory. The average

concentrations of the heavy metals were showed the following descending order: Fe> Zn > Cr >Mn> Pb> Al> Cu> Ni > Cd>Hg.

For irrigation purpose, average values of PH, HCO₃, SO₄, Ca, Pb, Cd, Cu, Fe, Ni, Al and Mn were less than the values of ²⁶. Other values may be more than or not detected in the ²⁶ standards (Table. 4).

Finally, the physico- chemical parameters of this study compared with many parameters in different years (Table 9). Some parameters ranges of the present study more than the most previous studies such as PH,DO, BOD, COD, CO₃ and Ca, while the others may be less than or not detected.

Assessment of Water Quality:

Tables 10 and11 and Figs. 3,4,5 and 6 illustrate the values of the WQI, HMPI, MPI and PI of Damietta Nile Branch water in Damietta Governorate. The WQI, HMPI, MPI and PI scores for Drinking water was computed using guidelines of ^{25,24}. Guidelines of ²⁶ were used to compute the WQI, HMPI, MPI and PI value for irrigation water.

1- Water Quality Index (WQI):

The parameters PH, TDS, Ec, DO, BOD, COD, CO₃, HCO₃, Cl, SO₄, Ca, Mg, NO₃, Hardness and alkalinity were used to calculation the WQI for drinking and irrigation water (Table. 10 and 11). For drinking water, the water quality was unsuitable in samples 1,2,3 and4 (these samples in the north near from the Mediterranean Sea). WQI were classified as poor in samples 5,6,7,8,12 and 13, while the water is good in the samples 9,10,11 and 14 (toward south) (Table 10, Fig. 3). For Irrigation water, the water quality was excellent in most samples (samples 6 to 14), good in samples 5, poor in samples 3 and 4 and the water was unsuitable for irrigation in samples 1 and 2 (near the Nile outlet to the sea) (Table 11, Fig. 3).

Table (9): Physical and chemical characteristics of Dameitta branch water during different time period compared with the present results

Parameters	[44]	[19]	[45]	[46]	Present results (Damietta Governorate)
Wat. Temp. 0C	15.9-32.3	17-33.6	16.1-27.5	17-35	
Trans. cm	35-90	70-150	45-200	50-150	
EC μ mohs/cm	500-44000	316-503	1007-69260	330-50200	370-22452
Salinity %	ND-30	-	9.87-42	ND-32	
TS mg/L	502-34300	186-454	710-41340	256-56302	215- 14369
pH	7.66-8.05	6.97-8.45	7.53-7.97	7.24-8.30	7.1 – 8.41
DO mg/L	6.07-7.23	6.60-9.60	3.03-10.37	5.5-11.0	6.2 – 12.5
BOD mg/L	3.45-8.45	1.20-3.50	1.67-7.70	1.5-4.0	2.2- 4.9
COD mg/L	6.92-10.12	2.8-8.8	9.47-24.67	8.0-26.0	11 - 28
CO ₃ ²⁻ mg/L	2.5-14.7	nil-5	nil-30.13	nil-10	7- 15
HCO ₃ ⁻ mg/L	155-174	96-156	115-150	150-220	122 - 200
Cl ⁻ mg/L	75-17835	11.34-36	38-24480	35-25500	33 - 8653
SO ₄ ²⁻ mg/L	137-2643	13.87-45.40	60-10970	30-3000	24 - 290
Ca ²⁺ mg/L	19.3-187.8	27-38	26-803	28-290	22 - 322
Mg ²⁺ mg/L	11-590	17.5-3016	20-1850	19-1300	18 - 421
NO ₂ - μ g/l	25-128	3.70-37.91	10-96	11-36	
NO ₃ - μ g/l	233-1056	11.19-89.75	27-386	30-100	15 - 44
NH ₃ mg/L	0.91-14.54	0.38-1.08	0.35-1.895	0.15-1.0	
PO ₄ ³⁻ μ g/l	166-259	11.24-9605	45-417	35-255	
TP μ g/l	338-574	48-283	290-904	250-440	
SiO ₃ ²⁻ mg/L	4.36-5.27	1.07-5.55	1.65-6.00	1.0-3.0	
T- Hardness mg/L					41 - 743
Alkalinity mg/L					25 - 126

2- Heavy Metals Pollution Index (HMPI)

Heavy metal pollution index (HMPI) is an effective method to evaluate the surface water quality. Where, HMPI values in drinking water were high heavy metal pollution (HHMP) in samples which near the Damietta Nile branch outlet (samples 1,2,3 and 4) and they were low heavy metal pollution (LHMP) in sample from 5 to 14 (Table 10, Fig. 4). For irrigation water, HMPI values were low heavy metal pollution (LHMP) in whole samples (Table 11 and Fig. 4).

3- Metal Pollution Index (MPI)

Metal pollution index is also an effective method to evaluation the quality water for drinking and irrigation purposes. In Table 10 and Figure 5 the MPI showed that all samples were non-potable water for drinking purpose. For irrigation, the samples were suitable for irrigation purpose in samples from 5 to 14, while the water in samples from 1 to 4 were high danger for irrigation purpose (Table 11 and Fig. 5).

4- Pollution Index (PI):

Pollution index is also an effective method to evaluation the quality water for drinking and irrigation purposes. The water quality according to PI was as following: Samples from 5 to 14 and sample 3 were classified as no effect, they slightly effect in samples 1 and 2 (Near outlet of Damietta Nile Branch) and strongly effect in sample 4 (Table 10 and Fig. 6). For irrigation water, all samples classified as no effected water by heavy metals (table 11 and Fig. 6).

Table.(10): Quality of Drinking Water of Damietta Nile Branch in Damietta Governorate

Samples	WQI	Quality of Drinking water	HMPI	Quality of Drinking water	MPI	Quality of Drinking water	PI	Quality of Drinking water
1	504	Unsuitable	124.8	HHMP	10.00	non-potable	1.08	Slightly affected
2	368	Unsuitable	130.4	HHMP	9.52	non-potable	1.02	Slightly affected
3	162	Unsuitable	106.9	HHMP	8.51	non-potable	0.85	No effect
4	123	Unsuitable	366.5	HHMP	30.4	non-potable	3.13	Strongly affected
5	60	Poor	77.8	LHMP	5.55	non-potable	0.57	No effect
6	56	Poor	70.01	LHMP	4.97	non-potable	0.59	No effect
7	56	Poor	78.00	LHMP	6.50	non-potable	0.58	No effect
8	59	Poor	45.11	LHMP	4.09	non-potable	0.37	No effect
9	44	Good	47.44	LHMP	3.52	non-potable	0.38	No effect
10	50	Good	62.1	LHMP	5.33	non-potable	0.57	No effect
11	44	Good	54.19	LHMP	4.33	non-potable	0.43	No effect
12	61	Poor	111.7	HHMP	9.05	non-potable	0.99	No effect
13	55	Poor	44.3	LHMP	3.18	non-potable	0.35	No effect
14	40	Good	41.9	LHMP	2.97	non-potable	0.33	No effect

HHMP: High heavy metal pollution

LHMP: Low heavy metal pollution

Table.(11): Quality of Irrigation Water of Damietta Nile Branch in Damietta Governorate

Samples	WQI	Quality of Irrigation water	HMPI	Quality of Irrigation water	MPI	Quality of Irrigation water	PI	Quality of Irrigation water
1	255	Unsuitable	12.85	LHMP	1.22	High danger	0.188	No effect
2	181	Unsuitable	11.54	LHMP	1.05	High danger	0.131	No effect
3	73	Poor	11.93	LHMP	1.07	High danger	0.133	No effect
4	55	Poor	16.08	LHMP	1.54	High danger	0.228	No effect
5	26	Good	4.78	LHMP	0.383	suitable	0.047	No effect
6	23	Excellent	7.52	LHMP	0.698	suitable	0.087	No effect
7	25	Excellent	5.35	LHMP	0.458	suitable	0.057	No effect
8	24	Excellent	3.96	LHMP	0.346	suitable	0.043	No effect
9	17	Excellent	7.10	LHMP	0.632	suitable	0.079	No effect
10	18	Excellent	7.08	LHMP	0.752	suitable	0.094	No effect
11	17	Excellent	4.5	LHMP	0.412	suitable	0.051	No effect
12	21	Excellent	8.21	LHMP	0.91	suitable	0.113	No effect
13	20	Excellent	9.15	LHMP	0.779	suitable	0.097	No effect
14	14	Excellent	8.71	LHMP	0.729	suitable	0.091	No effect




Fig. (3): WQI of Drinking and Irrigation water of Damietta Nile Branch in Damietta Governorate




Fig. (4): HMPI of Drinking and Irrigation water of Damietta Nile Branch in Damietta Governorate




Fig. (5): MPI of Drinking and Irrigation water of Damietta Nile Branch in Damietta Governorate




Fig. (6): PI of Drinking and Irrigation water of Damietta Nile Branch in Damietta Governorate

Conclusions:

The parameters (PH, TDS, Ec, DO, BOD, COD, CO₃, HCO₃, Cl, SO₄, Ca, Mg, NO₃, Hardness and alkalinity) and heavy metals concentration used for evaluation of water for drinking and irrigation purposes. Most of water suitable for irrigation, while some of water suitable for drinking. The water near the Damietta Nile Branch Outlet is unsuitable for drinking and irrigation, but the water in the south of study area suitable for all purposes.

References:

1. Wahaab, R.A. and Badawy, M. I. (2004): Water Quality Assessment of the River Nile System: An Overview. *Biomedical And Environmental Sciences* 17, 87-100 (2004) .
2. World Bank (2005) Country Environmental Analysis (1992-2002), Arab Republic of Egypt, Water and Environment Department. The Middle East and North Africa Region. World Bank.
3. Chapman, D. and Chapman, D. E (1996) "Water Quality Assessments. A Guide to the Use of Biota, Sediments and Water in Environmental Monitoring," 2nd Edition, Chapman & Hall, London, 1996.
4. Mapfumo, E. Willms W. and Chanasyk, D. (2002) "Water Quality of Surface Runoff from Grazed Fescue Grassland Watershed in Alberta," *Water Quality Research Journal of Canada*, Vol. 37, No. 3, 2002, pp. 543-562.
5. ELEWA, H. H. 2010. Potentialities of Water Resources Pollution of the Nile River Delta, Egypt. *The Open Hydrology Journal* 4 : 1- 13.
6. Fernandez, G., Chescheir, G. M Skaggs R. W. and Amatya, D. M. (2002) "WATGIS: A GIS-Based Lumped Parameter Water Quality Model," *Transactions of the ASAE*, Vol. 45, No. 3, 2002, pp. 593-600.
7. NWRC,(2000), National Water Quality and Availability Management: National Water Quality Monitoring Component 100. Final Report, National Water Research Center (NWRC) and Agriculture and Agri-Food Canada (PFRA), Egypt.
8. Badawy, M.I. (2013). Water Quality of Drinking Water Resources and Formation of Disinfection by-products. Egyptian-German Workshop on Sustainable Water Technologies (SusWaTec Workshop) 18.-20th February 2013-Cairo, Egypt. National Research Centre, Water Pollution Research Department Dokki, Cairo, Egypt
9. Sundaray, S. K., Panda, U. C., Nayak, B. B., & Bhatta, D. (2006). Multivariate statistical techniques for the evaluation of spatial and temporal variation in water quality of Mahanadi river-estuarine system (India), A case study. *Environmental Geochemistry and Health*, 28(4), 317-330.

- <http://dx.doi.org/10.1007/s10653-005-9001-5>.
10. Akoto, O., Bruce, T. N., &Darko, G. (2008). Heavy metals pollution profiles in streams serving the Owabi reservoir. *African Journal of Environmental Science and Technology*, 2(11), 354-359.
 11. Amadi, A. N., & Olasehinde, P. I. (2010). Application of remote sensing techniques in hydrogeological mapping of parts of Bosso Area, Minna, North-Central Nigeria. *International Journal of Physical Sciences*, 5(9), 1465-1474.
 12. Gueu, S., Yao, B., Adouby, K., & Ado, G. (2007). Kinetics and thermodynamics study of lead adsorption on to activated carbons from coconut and seed hull of the palm tree. *International Journal of Environmental Sciences and Technology*, 4(1), 11-17.
 13. Lee, C. L., Li, X. D., Zhang, G., Li, J., Ding, A. J., & Wang, T. (2007). Heavy metals and Pb isotopic composition of aerosols in urban and suburban areas of Hong Kong and Guangzhou, South China Evidence of the long-range transport of air contaminants. *Environmental Pollution*, 41(1), 432-447.
 14. Adams, R. H., Guzmán-Osorio, F. J., & Zavala, C. J. (2008). Water repellency in oil contaminated sandy and clayey soils. *International Journal of Environmental Science and Technology*, 5(4), 445-454.
 15. Vinodhini, R., & Narayanan, M. (2008). Bioaccumulation of heavy metals in organs of fresh water fish *Cyprinus carpio* (Common carp). *International Journal of Environmental Science and Technology*, 5(2), 179-182.
 16. Suthar, S., & Singh, S. (2008). Vermi composting of domestic waste by using two epigeic earthworms (*Perionyx excavates* and *Perionyx sansibaricus*). *International Journal of Environmental Science and Technology*, 5(1), 99-106.
 17. Aktar, M. W., Paramasivam. M., Ganguly, M., Purkait, S., & Sengupta, D. (2010). Assessment and occurrence of various heavy metals in surface water of Ganga river around Kolkata: a study for toxicity and ecological impact. *Environmental Monitoring and Assessment*, 160(2), 207-213. <http://dx.doi.org/10.1007/s10661-008-0688-5>
 18. Abdo, M. H. (2004). Environmental studies on the River Nile at Damietta branch region, Egypt. *J. Egypt. Acad. Soc. Environ. Develop., (D- Environmental Studies)*, 5 (2): 85 - 104.
 19. Al-Afify, A. D. G. (2006). Biochemical studies on River Nile pollution. M.Sc. Thesis, Biochemistry Department Fac. Agric., Cairo University, Giza, Egypt. 152pp.
 20. APRP. (Agricultural Policy Reform Program, Water Policy Program), 2002. Survey of Nile System Pollution Sources. Agricultural Policy Reform Program (APRP), Ministry of Water Resources and Irrigation, Report No. 64, Cairo, Egypt
 21. El-Gammal, M.I., Ali, R.R and Eissa, R. (2014): Land use assessment of barren areas in Damietta Governorate, Egypt using remote sensing. *Egyptian journal of basic and applied sciences*, I (2014) 151-160.
 22. Adams, V. D (1990) *Water and Wastewater Examination Manual*, Published by Taylor and Francis (1990), ISBN 10: 0873711998 ISBN 13: 9780873711999
 23. APHA, (2005): *Standards methods for examination of water and wastewater*. American Public Health Association, Washington, D.C.
 24. WHO, (2011) *Guidelines for Drinking Water Quality*, Water Health Organization, Fourth Edition, pp. 219- 443.
 25. Egyptian drinking water quality standards, 2007. Ministry of Health, Population Decision number (458).
 26. (FAO), (1994): *The state of food and agriculture*. FAO Agriculture Series, No. 27. ISSN 0081-4539, Rome, 1994.
 27. Abbasi, T. and Abbasi, S.A. (2012) *Water Quality Indices*, Elsevier B.V., pp. 19-28.
 28. Brown, R.M, N.J. Mc Clelland, R.A. Deiniger, and M.F.A. O' Connor (1972). Water quality index – crossing the physical barrier (Jenkins, S H ed) *Proc. Intl. Conf. on water poll. Res. Jerusalem* 6, 787 – 797.
 29. El-Sherbini A. and El-Moattassem M. (1994). *River Nile Water Quality Index during High and Low Flow Conditions*. National Conference on the River Nile, Assiut Center for Environmental Studies.
 30. Mohanta, B.K. and Patra, A.K. (2000). Studies in the water quality index of river Sanamachhakananda at Keonjargarh, Orissa, India. *Pollut. Res.*, 19, 377-385.
 31. Dwivedi S. L. and V. Pathak, (2007). A Preliminary Assignment of Water Quality Index to Mandakini River, Chitrakoot, *Indian Journal of Environmental Protection*, Vol. 27, No. 11, pp. 1036-1038.
 32. Simoes F. S., A. B. Moreira, M. C. Bisinoti, S. M. N. Gimenez and M. J. S. Yabe, (2008). Water Quality Index as a Simple Indicator of Aquaculture Effects on Aquatic Bodies, *Ecological Indicators*, Vol. 8, No. 5, pp. 476-484.

33. Chatterjee, C. and Raziuddin, M. (2002): Determination of water quality index of a degraded river in Asanol Industrial area, Raniganj, Burdwan, West Bengal, *Nature, Environment and Pollution Technology*, 1 (2), 2002, pp 181-189.
34. Bakan, G. , Özkoç, H.B., Tülek, S., Cüce, H.(2010):Integrated environmental quality assessment of the Ki{dotless}zi{dotless}li{dotless}rmak River and its coastal environment, OndokuzMayis University, Engineering Faculty, Department of Environmental Engineering, 55139, Samsun, Turkey. *Turk. J. Fisheries Aquat. Sci.* 10, 453-462.
35. Tamasi, G., &Cini, R. (2004).Heavy metals in drinking waters from Mount Amiata. Possible risks from arsenic for public health in the province of Siena. *Science of the Total Environment*, 327, 41-51. <http://dx.doi.org/10.1016/j.scitotenv.2003.10.011>
36. Mohan, S.V., Nithila, P., and Reddy, S.J., 1996. Estimation of heavy metals in drinking water and development of heavy metal pollution index: *Journal of Environmental Science & Health Part A*, v. 31, p. 283-289.
37. Prasad, B., &Kumari, S. (2008). Heavy metal pollution index of ground water of an abandoned open cast mine filled of the water quality of River Adyar, India. *Bulletin of Environmental Contamination and Toxicology*, 82(2), 211-217.
38. Chon HT, Ahn JS, Jung MC (1991) Environmental contamination of toxic heavy metals in the vicinity of some Au–Ag mines in Korea, *Proc. of the 4th. Biennial SGA Meeting.Truku, Finland*, p 891.
39. Kim KW, Lee HK, Yoo BC (1998) The environmental impact of gold mines in the Yugu- Kwangcheon Au-Ag metallogenic province Republic of Korea. *Environ SciTechnol* 19:291–298.
40. Emoyan O.O, Ogban FE, Akarah E (2005) Evaluation of Heavy metals loading of River Ijana. *Nigeria J ApplSci Environ Manag* 10(2):121–127.
41. Odukoya AM, Abimbola AF (2010) Contamination assessment of surface and groundwater within and around two dumpsites. *Int J Environ Sci Tech* 7(2):367–376
42. Lee JS, Chon HT, Kim JS, Kim KW, Moon HS (1998) Enrichment of potentially toxic elements in areas underlain by black shales and slates in Korea. *Environ GeochemHlth* 20(30):135–147
43. Caerio, S., Costa, M. H., Ramos, T. B., Fernandes, F., Silveira, N., Coimbra, A., Painho, M. (2005). Assessing heavy metal contamination in Sado Estuary sediment: An index analysis approach. *Ecological Indicators*, 5, 155-169.
44. Sayed, M. F. (1998).Evaluation of pollution on Mugilspecies in Damietta branch of the River Nile between Faraskour Barrage and Ras El-Bar outlet. M. Sc. Thesis, Fac. Sci., Helwan Univ., Egypt.
45. Mohamed, M. M. (2010). Ecological studies on the effect of Faraskour Dam on the water quality of Damietta branch of the River Nile. M. Sc. Thesis, Fac. of Sci Al-Azhar Univ.
46. Abdo M. H (2010) Environmental and water quality evaluation of Damietta branch, River Nile, Egypt. *African J. Biol. Sci.*, 6 (2): 143-158. ISSN 1687-4870.
