



## Biochemical and Clinopathological studies in Common Carp (*Cyprinus Carpio*, L.) Fish exposed to pollution by Heavy Metals

Mona S. Zaki <sup>1\*</sup>, Olfat M. Fawzi <sup>2</sup> and Refat A. Youssef <sup>3</sup>

<sup>1</sup>Department of Hydrobiology, National Research Center, Egypt

<sup>2</sup>Biochemistry Department, National Research Center, Egypt

<sup>3</sup>Soil and water use Department, National Research Center, Egypt

**Abstract :** Heavy metals are persistent contaminants in the environment causing serious illness in fish, animals and human. Lead represents the main toxic element in nature. Lead has a tendency to accumulate in tissue and organs of exposed fish. The present study aimed to investigate the effect of lead pollution on fish with special reference to the hematological, immunological, serum biochemical parameters, where fifty healthy Common Carp fish were divided into 3 groups. Fish of gp1 served as a control. Fish of gp. 2 & 3 were used for the determination of acute lethal concentration dose and the pathological effect of lead on the exposed Common Carp. Blood samples were collected to obtain serum for biochemical studies and heparinized blood for hematological investigations. RBCs, Hb, HCT, and MCHC showed significant elevations, the serum GPT and GOT were increased significantly. L.D.H, glucose and cortisol were elevated, while serum cholesterol concentration was reduced significantly in high tem30°C.

**Key words :** Lead pollution, Common Carp, Biochemical changes.

### 1. Introduction

Lead occurs naturally in the environment. However, most lead concentrations that are found in the environment are a result of human activities. Due to the application of lead in gasoline, an unnatural lead-cycle has consisted. In car engines lead is burned, so that lead salts (chlorines, bromines, oxides) will originate. These lead salts enter the environment through the exhausts of cars. The larger particles will drop to the ground immediately and pollute soils or surface waters, the smaller particles will travel long distances through air and remain in the atmosphere. Part of this lead will fall back on earth when it is raining. This lead-cycle caused by human production is much more extended than the natural lead-cycle, and has caused lead pollution to be a worldwide issue<sup>1</sup>.

The high level of lead could be also due to the industrial discharges from superphosphate factories, traffics of high way or motor vehicles as well as the extensive use of agrochemicals such as fertilizers, pesticides and growth promoters<sup>2</sup>.

Lead can enter the human body through uptake of food (65%), Water (20%) and air (15%) and cause several unwanted effects, such as: Disruption the biosynthesis of haemoglobin and anaemia, a rise in blood pressure, kidney damage and Miscarriages and subtle abortions, disruption of nervous systems, brain damage, declined fertility of men through sperm damage and diminished learning abilities of children and behavioural disruptions of children, such as aggression, impulsive behavior and hyperactivity. Lead can enter a fetus

through the placenta of the mother. Because of this, it can cause serious damage to the nervous system and the brains of unborn children<sup>3</sup>.

Industrial and agricultural discharges are considered the primary source of metal poisoning to fish in Egypt<sup>4</sup>. Lead has a tendency to accumulate in tissue and organs of exposed fish resulting in hepatic and renal dysfunction with growth retardation<sup>5</sup>. Consequently, it could induce alterations in hematological and serum biochemical parameters<sup>6</sup> as well as pathological changes in most body organs<sup>7</sup>.

The present study aimed to investigate the effect of lead pollution on fish with special reference to the haematological, immunological, serum biochemical parameters.

## **Material and Methods:**

### **1- Fish:**

Fifty healthy Common Carp fish of both sexes and  $150 \pm 50$  gm body weight, were obtained alive and transported immediately to the laboratory. They were kept in 5 glass aquaria (100 X 30 X 50 cm) that provided daily with a tap water and continuously with filtered air. The water temperature was adjusted at 30°C along the period of experiment using thermostatic heater. The fish were fed a balanced ration daily using the formula suggested by *Ahmed and Matty*, 1988 [8]. Fish were kept under observation for 2 weeks.

Fish were divided into 3 groups (gps). Fish of gp1 (10) served as a control with no treatment. Fish of gp. 2 & 3 (20, each) were used for the determination of acute lethal concentration dose (LD<sub>50</sub> /72 hr, gp2) and to investigate the pathological effect of lead on the exposed fish (gp3).

### **2-Experiments:**

#### **A- Determination of acute lethal concentration dose:**

To determine lethal concentration dose, fish of gp. 2 were subdivided into 5 equal subgroups. Subgroup 1 served as a control. Other 4 subgroups exposed to 35, 75, 150 and 300 mg/L of lead acetate; respectively. Each dose was dissolved in the distal water of each aquarium. The number of dead fish was recorded within 72 hrs post-exposure and the acute lethal concentration dose was calculated according to the formula of Brown,<sup>9</sup>.

#### **B- long term exposure:**

Fish of gp3 were exposed to 1/100 of LD<sub>50</sub> /72 hr (1 mg/L) of lead acetate for 2 weeks according to Taylor et al.,<sup>10</sup>. The excreta were removed regularly and the water was replaced within 4 days interval. Fish were kept under observation along the 14 days of exposure.

### **3- Sampling:**

Blood sample were collected from the caudal vein after 3, 7, 14 days of exposure, part of blood was left to clot and then centrifuged at 3000 r.p.m. to obtain serum for biochemical studies, the other part was heparinized for hematological investigations using the methods of Drabkin,<sup>11</sup>.

### **4-Haematological examinations:-**

The erythrocytic indices (RBCs, Hb, Hct & MCHC) were estimated according to Schalm<sup>12</sup>.

### **5-Serum biochemical analysis:**

Kits Biomerieux France were used for the determination of serum glutamic pyruvic transaminase (GPT) and glutamic oxaloacetic transaminase (GOT), lactate dehydrogenase (LDH), alkaline phosphatase (AP), serum glucose, serum cholestrol, and total protein. Serum cortisol hormone was analyzed by means of a gemmacoat 125-cortisol radio-immunassay kit (Diagnostic corporation, USA). Serum Ig M was also measured according to Fuda, et al.,<sup>13</sup>.

## 6-Statistical analysis:

The obtained data were statistically analyzed according to Snedecor and Cochran, <sup>14</sup> by T test.

## 3. Results

### A-Determination of acute lethal concentration dose:

Experiment 1 revealed that, the acute lethal concentration dose was 150 mg/L during the 1<sup>st</sup> 72 hrs post-exposure.

### B- Long term exposure:

The effect of lead acetate exposure on RBC's count, Hb level, Hct and MCHC values of exposed tilapia fish were recorded in Table (1). Polycythemia was observed on the 14 day ( $p < 0.01$ ). Blood Hb, Hct, and MCHC showed a significant elevation by 14 days of experiment.

**Table 1: Effect of lead on some haematological parameters of Common Carp fish along the period of experiment (Mean±S.E.)**

Parameter	3days		7days		14days	
	Control	Exp.	Control	Exp.	Control	Exp.
RBCs $10^6/\text{mm}^3$	$3.1 \pm 0.22$	$3.1 \pm 0.51$	$2.9 \pm 0.33$	$3.9 \pm 0.74$	$3.4 \pm 0.59$	$4.5 \pm 0.68^*$
HB g/dl	$7.00 \pm 0.44$	$8.1 \pm 0.18$	$6.9 \pm 0.13$	$8.4 \pm 0.60$	$7.1 \pm 0.20$	$9.0 \pm 0.54^{**}$
H.Ct%	$18.80 \pm 1.00$	$21.9 \pm 1.20$	$18.95 \pm 0.25$	$26.4 \pm 1.54$	$20.7 \pm 1.64$	$27.7 \pm 1.84^{**}$
MCHc%	$31.70 \pm 1.60$	$32.8 \pm 1.30$	$30.52 \pm 0.74$	$35.52 \pm 1.06$	$30.3 \pm 1.22$	$39.6 \pm 1.07^{**}$

Exp: experimental \* Significant at  $p < 0.01$ . \*\* Non-significant

**Table 2: Effect of lead on the serum biochemical parameters of Common Carp fish along the period of experiment (Mean±S.E.)**

Parameter	3days		7days		14days	
	Control	Exp.	Control	Exp.	Control	Exp.
SGPT (I.U/L)	$26.2 \pm 1.1$	$49.3 \pm 0.34$	$28.3 \pm 0.74$	$41.5 \pm 1.09$	$30.0 \pm 1.00$	$46.9 \pm 2.18^*$
SGOt (I.U/L)	$37.3 \pm 1.8$	$39.7 \pm 2.5$	$39.8 \pm 1.22$	$46.9 \pm 2.18$	$37.32 \pm 1.1$	$50.40 \pm 3.04^{**}$
L.D.H (I.U/L)	$185 \pm 4.3$	$185 \pm 3.04$	$185 \pm 4.04$	$185 \pm 4.23$	$190 \pm 3.00$	$193 \pm 4.28^*$
A.L.P (U/L)	$3.50 \pm 1.1$	$2.8 \pm 1.04$	$3.9 \pm 1.54$	$4.5 \pm 1.40$	$2.32 \pm 1.20$	$5.64 \pm 2.15^{**}$
Glucose (mg / dl)	$27.24 \pm 1.5$	$31.68 \pm 1.1$	$27.82 \pm 1.1$	$40.20 \pm 2.0$	$28.64 \pm 1.1$	$58.8 \pm 2.38^{**}$
Total protein (g/dl)	$2.07 \pm 0.22$	$2.22 \pm 0.11$	$2.40 \pm 0.54$	$2.8 \pm 0.24$	$2.4 \pm 0.45$	$4.59 \pm 0.54^*$
Cholesterol (ng/dl)	$116.4 \pm 3.0$	$122.6 \pm 2.3$	$122 \pm 2.6$	$168 \pm 3.03$	$117.9 \pm 2.1$	$190.4 \pm 4.0^*$
Ig. M (ng/ml)	$1.6 \pm 0.10$	$1.45 \pm 0.20$	$1.61 \pm 0.32$	$1.23 \pm 0.72$	$1.55 \pm 0.11$	$0.3 \pm 0.065^*$
Cortisol (ng/ml)	$0.75 \pm 0.21$	$1.13 \pm 0.05$	$0.89 \pm 1.03$	$1.67 \pm 1.00$	$0.75 \pm 0.52$	$1.24 \pm 1.24^*$

Exp: experimental \* Significant at  $p < 0.01$ . \*\* Non-significant

Table (2) revealed the changes of some biochemical constituents in the blood of tilapia fish due to lead acetate exposure. The obtained data revealed that serum GPT activity increased significantly by 14 days of exposure. A significant elevation in serum GOT activity was also observed on the 14<sup>th</sup> day ( $p < 0.01$ ). L.D.H serum activity was elevated along the whole period of experiment especially on the day 14<sup>th</sup>. Hyperglycemia was constant findings from the beginning of the experiment until the end of the experiment. Serum cholesterol concentration was increased, on the 3<sup>rd</sup> day and the 7<sup>th</sup> day and was reduced significantly, on the day 14<sup>th</sup> day. Cortisol hormone was elevated along the whole period of experiment especially on day 14<sup>th</sup>.

## Discussion:

Regarding the impact of lead on the hematological profile of Common Carp, polycythemia accompanied by elevated hemoglobin level, HCT value and MCHC were observed. Similar findings were reported by Mckim *et al.*<sup>15</sup>, Hilmy *et al.*<sup>16</sup> and Taylor *et al.*<sup>10</sup> recorded polycythemia in rasy barb. But in contrary to our finding Hb level and MCHC were reduced in Common Carp exposed to copper<sup>17</sup>. The increased RBCs count may be due to stimulation of erythropoietin by elevated demands for O<sub>2</sub> or CO<sub>2</sub> transport as a result of increased metabolic activity or distraction of gill membranes causing faulty gaseous exchange. The increase Hb content could be explained as a process where the body tries to replace the oxidized denatured Hb<sup>18</sup>. The increase of HCT value and MCHC may be attributed to swelling of RBCs due to increased CO<sub>2</sub> in blood, hypoxia or stressful procedures<sup>19,20</sup>.

Exposure of Common Carp to sublethal concentration (1.5 mg/L) of lead acetate for 14 days resulted in a marked increase in the activities of serum GPT, GOT, LDH and ALP. The present findings agree with our microscopic findings, which revealed a marked degeneration and necrosis of hepatocytes as the elevation in transaminases activities may be attributed to the liver injury<sup>21</sup>.

Serum cholesterol level, in the present study, showed a significant reduction that could be due to greater level of utilization of cholesterol during corticosteroidogenesis, as it is the precursor for steroid hormones [22]. In addition, they reported a rise in the blood protein resulted in a high density of lipoprotein in the serum and was suggested to be the cause of hypocholesterolemia in exposed fish.

Our results showed similar findings as that of Gill *et al.*,<sup>6</sup> and Snieszko<sup>23</sup>, who reported that, exposure of fish to lead had no significant increase on blood glucose of *salmo gairnei*. The blood glucose level reflected the changes in carbohydrate metabolism under hypoxia and stress conditions. Rise of glucose level indicated the presence of stressful stimuli eliciting rapid secretion of both glucocorticoids and catecholamines from the adrenal tissue and accompanied by cortisol elevation<sup>24</sup>. Concerning serum protein level, a significant increase was noted 14 days post exposure to lead. The elevated protein concentration may be due to the induction of protein synthesis in liver.

The serum Ig. M was determined to find out information about fish immune system which was previously investigated in different species by many authors as Fuda *et al.*,<sup>13</sup> and O'Neill<sup>25</sup>, in this work, the purified Ig. M was revealed a single perception against specific polyvalent antiserum to tilapia fish Ig, similar results was obtained by Bagee *et al.*,<sup>26</sup> who found that, *Coho salmon* Ig was detected by specific anti Ig 14. Our study revealed a significant decrease in Ig. M level in fish exposed to lead pollution if compared within control groups. Anderson *et al.*,<sup>27</sup> found a relation between cortisol and IgM as when cortisol increased IgM decrease. The significant increase in cortisol level in fish exposed to lead could be attributed to stress factors and the intoxication of fish<sup>28</sup>.

## References

1. Dowidar M., Abdel-Magid S. and Salem S. 2001: Biochemical effect of lead and cadmium on glutathion peroxidase superoxide dismutase activity, copper and selenium level in rat. 2<sup>nd</sup> Int. sc. Conf., Fac. vet. Med., Mansura univ.
2. Tolba K., El-Neklawy G. and Niazi Z. 1994: Lead residue in tissue of slaughtered cattle kept at agriculture and industrial areas. J. Egypt. Vet. Med. Ass., 54 (2) 199.
3. Jarup L. 2003: Hazards of heavy metals contamination. Br. Med. Bull; 68: 167-82
4. El-Nabawi A., Heinzow B. and Kruse H. 1987: Cd, Cu, Pb, Hg and Zn in As fish from Alexandria region of Egypt. Bull. Environ. Contam. Toxicol., 39, 889-897.
5. Haneef S., Swarap D. and Dwivedi S. 1998: Effects of concurrent exposure to lead and cadmium on renal function in goats. Indian vet. Research, 28, 3, 257 – 261.
6. Gill S.; Tewari H. and Ponde J. 1991: Effect of water born copper and lead on the peripheral blood in the *rosy barb, barbush*. Bull. Environ. Contam. Toxicol., 46, 606-612.
7. Ghalab M. 1997: Clinicopathological studies on fish exposed to some environmental pollution in El-Manzala lake. Ph. D. thesis, Fac. Vet. Med., Suez Canal Univ.
8. Ahmed T. and Matty A. 1989: The effect of feeding antibiotic on growth and body composition of carp (*Cyprinus carpio*). Aquaculture, 77, 211.

9. Brown V. 1980: Acute toxicity in theory and practice with special reference to the toxicology of pesticides. Wiley in science publication. John Willey and Sons, Chichester.
10. Taylor D.; Maddack B. and Murce G. 1985: The acute toxicity of mine grey list *metals* e.g arsenic, chromium, copper, lead, nickel, vanadium, and zinc marine fish species *lob limanda* and *grey mullet*. *Aqua. Toxicol.*, 6, 3. 135-145.
11. D. Drabkin 1949: Standardization of hemoglobin measurements. *Am. J. Med. Sci.*, 217-710.
12. Schalm O. 1986: Schalm's Veterinary Hematology. 4<sup>th</sup> Edition 524.
13. Fuda H., Sayano K., Yamaji F. and Haraj A. 1991: Serum immunoglobulin M (IgM) during early development of *masu salmon* on *corhyrchus masu*. *Comp. Biochem. Physiol.*, 99, 637.
14. Snedecor F. and Cochran S. 1969: Statistical analysis. Iowa State Univ. Press. Iowa, USA.
15. McKim J.; Christensen G.; Hunt E. 1970: Changes in the blood of brook trout *salsvlinus* after short term and long term exposure to copper. *Fish Des Bd Con.*, 27, 1883-1889.
16. Hilmy M.; Lemke A.; Jaicb P. 1979: Haematological changes in Kuwait mullet, *liza maeralepis* (smith induces by heavy metals). *Indian Mar. Sci.* 8, 278-281.
17. El-Domiaty N. 1987: Stress response of juvenile *Clarias lazera* elicited by copp. *Comp. Bioch. Physiol.*, 88, 259-262.
18. Cyria P., Antony H., and Nonbisor P. 1989: Haemoglobin and haematocrit values in the fish *Oreochromis mossambicus*, after short term exposure to copper and lead. *Bull. Environ. Contam. Toxicol.* 43, 315-320
19. Ellis A. 1981: Stress and the modulation of defence mechanisms in fish in pickering A.D. Stress and Fish Academic Press, New York, 147-169.
20. Nemesok J. and Boross L. 1999: Comparative studies on the sensitivity of different fish species to metal pollution. *Hoto Biol. Hung* 33, 27-27.
21. Kristaffevsson R. and Okari P. 1974: Effect of sublethal concentration of phenol on plasma enzyme activities in *pike* in brakish water. *Ann. Zool. Fennici*, 11, 220-223.
22. Ferranda M. and Andrew M. 1991: Effect of lindane on the blood of fish water. *Bull. Environ. Contam. Toxicol.*, 47, 465 – 470.
23. Sniezko S. 1974: The effect of environmental stress on outbreaks of infectious diseases in fish. *Journal of Fish Biol.*, 6, 197 – 208.
24. Mareaud M.; Mareaud F. and Donlds E. 1977: Primary and secondary effects of stress in fish, some new data with a general review. *Trans. Amer. Fish Sac.*, 106, 201.
25. O'Neill J. 2001: The humoral immune response of *salmo trutta* exposed to heavy metals. *J. Biol. Chem.*, 201, 706-721.
26. Bagee M.; Fuda H., Hara H.; Kawamura H. and Yamauchi H. 1993: Changes in serum immunoglobulin M. (IgM) concentrations during early development of *chum salmon* as determined by sensitive Elisa technique. *Comp. Biochem. Physiol.*, 106, 69.
27. Anderson D., Roberson B. and Dixon O. 1982: Immunosuppression induced by corticosteroid or an alkylating agent in *Rainbow trout*. *Dev. Comp. Immunol. Suppl.*, 2, 197.
28. Aly S. M., Zaki M. S. and El-Genaidy H. M. 2003: Pathological, biochemical, haematological and hormonal changes in catfish exposed to lead pollution, *J. Egypt Vet. Med. Assoc.*, 63: 331-342

\*\*\*\*\*