



PharmTech

International Journal of PharmTech Research

CODEN (USA): IJPRIF, ISSN: 0974-4304, ISSN(Online): 2455-9563
Vol.9, No.11, pp 54-57, 2016

Identification of Risk Factors caused neonatal Sepsis of Premature Neonatusin Adam Malik General Hospital Center

Yettrie Bess Congencya Simarmata^{1*}, Urip Harahap¹, GuslihanDasa Djipta²

¹Faculty of Pharmacy, University of North Sumatera, Medan, Indonesia

²Faculty of Medicine, University of North Sumatera, Medan, Indonesia

Abstract :Until now, neonatal sepsis remains a major cause of mortality and morbidity of newborns. In developing countries, most of neonates treated with problems related to the infection. Giving TPN and utilizing unaseptic catheter will cause bacterial contamination in neonates. Neonatal sepsis is an infection that occurs in infants in the first 28 days after birth. Neonaturom Sepsis is categorized as early-onset sepsis (early onset) and late onset sepsis (late onset). This research aims to know the process of TPN and utilizing catheters as the biggest factor causing neonatal sepsis in premature neonates at the Haji Adam Malik General Hospital. This study is observational research which has been done on hospitalizedrespective, in Division of Perinatology General Hospital Haji Adam Malik in November 2014 – April 2015.

Keywords: neonatal sepsis, preterm neonates, risk factors.

Yettrie B CongencyaSimarmata *et al*/International Journal of PharmTech Research, 2016,9(11): 54-57.54
