

Suppressive effects of medicinal plants and their derivatives on inflammasome complex: a systematic review

Mohammad Reza Mahmoudian Sani^{1, 2}, Majid Asadi-Samani^{3*},
Ali Saeedi-Boroujeni⁴, Mehdi Banitalebi-Dehkordi², Mahmoud Bahmani⁵

¹Research Center for Molecular Medicine, Hamedan University of Medical Sciences, Hamedan, Iran;

²Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran;

³Student Research Committee, Shahrekord University of Medical Sciences, Shahrekord, Iran;

⁴Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran;

⁵Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran.

Abstract: Inflammasome activation is mediated by (NOD)-like receptors (NLR) proteins that respond to stimuli. Among NLRs, NACHT- LRR and PYD domains-containing protein 3 (NLRP3) senses the widest array of stimuli. NLRP3 inflammasome has an important role in the development of many inflammation disorders. Regarding the significance of inflammatory diseases, and the necessity of preventing and treating these diseases, the aim of this review article is to report medicinal plants and their nature-based derivatives that are effective on suppression of inflammasome complex. Web of Science and PubMed databases were searched using the Endnote software for the publications about the role herbal medicine in inflammasome activation pathways from 2000 to February 2016. *Sophora flavescens*, *Lycium barbarum*, *Impatiens textori* Miq., *Syneilesis palmata* (Thunb.), *Aloe vera*, citral (3,7-dimethyl-2,6-octadienal), celastrol, sulforaphane, schisandrin, resveratrol, dehydroniciferyl alcohol (DHCA), luteoloside, Pulsatilla decoction, and Wuling San have been reported to suppression function of inflammasome. Medicinal plants and their derivatives can be useful for inflammation related disorders by suppress NLRP3 inflammasome activation. However, they should be investigated in clinical trials to help to prevent and treatment of inflammatory diseases.

Keywords: Inflammasome, Inflammation, Medicinal plants, Phytochemicals, Herbal drugs.

Introduction

Inflammasomes are a group of large caspase-1-activating protein complexes. Inflammasomes response to evocation of innate immunity and production of pro-inflammatory cytokines such as LRP1, NLRP3, NLRC4, AIM2, and NLRP6 [1, 2]. They sense sensepathogen-associated molecular patterns (PAMPs) in the cytosol as well as damage-associated molecular patterns (DAMPs) [3, 4]. The inflammasomes play a role in cellular functions such as apoptosis and pyroptosis [5]. Explanation of inflammasome pathways with understanding of

how they contribute to autoimmune and autoinflammatory disorders alike might open up the horizon for the development of new therapies for these diseases [6]. In inflammasome complex, NLRs are activated by structural changes in them and bonding with ASC through pyrin domains. Furthermore, the adaptor protein ASC bonds with pro-caspase-1 through CARD domains. This big complex, called inflammasome, enables caspase-1 to self-activate. Then, caspase-1 is activated and converts inactive form of IL-1 β (pro-IL-1 β) or IL-18 into active form to prepare it for secretion and biological activity [7, 8]. Furthermore previous studies showed that inflammasome activation is mediated by ROS. Actually blockade of ROS production involving the use of chemical scavengers of ROS, and pharmacological inhibitors of NADPH oxidase inhibit NLRP3 inflammasome activation in response to a wide range of stimuli [9-11]. NLRP3 has been shown to play a central role in the induction of obesity and insulin resistance [12].

Normal activation of NLRP3 inflammasome contributes to host defense, but several studies suggest that excessive activation leads to the development of a number of inflammatory diseases [13]. Mutations of the human NLRP3 gene are associated with autoinflammatory syndrome, which is characterized by recurrent bouts of fever with debilitating local and systemic inflammation [14]. In mice, genetic ablation of NLRP3 or caspase-1 protects against obesity-associated inflammation and insulin resistance with reduced IL-1 in adipose tissue [15-17]. Activation of the NLRP3 inflammasome must be tightly regulated, and identification of anti-inflammatory agents that selectively target it would be very important. In this regards, suppression of NLRP3 inflammasome activation can be a new approach for prevention and treatment of inflammatory diseases.

Recently, medicinal plants have been attracting great attention and researchers have made efforts to discover their effects on cells, animals and human [18-25]. Numerous studies have confirmed the positive impact of medicinal plants and their derivatives, according to both folk and modern medicine, on prevention and treatment of various diseases [26-40]. Given the significance of inflammatory diseases, and the necessity of preventing and treating these diseases, the aim of this review article is to report medicinal plants and their nature-based derivatives that are effective on suppression of inflammasome complexes.

In this regards, Web of Science and PubMed databases were searched for the publications about the role of herbal medicine in suppression of inflammasome activation pathways 2000 and February 2016 using the EndNote software. The used search terms were inflammasome and medicinal plant or herb or herbal medicine or natural compound or phytochemical or herbal drugs and inflammasome. Each database was searched independently. The articles retrieved from both databases were analyzed once. Abstracts were reviewed based on predefined inclusion and exclusion criteria. When necessary, full texts were retrieved to assess study eligibility. Only the articles directly addressing the effect of the medicinal plants and their derivatives were selected and analyzed.

29 articles had investigated the role of the medicinal plants and their derivatives in regulating inflammasomes. *Sophora flavescens*, *Lycium barbarum*, *Impatiens textori* Miq., *Syneilesis palmata* (Thunb.), *Aloe vera*, citral (3,7-dimethyl-2,6-octadienal), celastrol, sulforaphane, schisandrin, resveratrol, dehydrodiconiferyl alcohol (DHCA), luteoloside, Pulsatilla decoction, and Wuling San have been reported to suppression of the function of inflammasome. Table 1 gives further details.

Sophora flavescens

S. flavescens as a traditional herbal medicine used to reduce inflammation, blocked the phosphorylation of IKK α/β , key upstream kinases involved in the degradation of I κ B α , and the cleavage of caspase-1. *S. flavescens* exerts its anti-inflammatory effects by blocking *P. aeruginosa*-mediated NF- κ B/inflammasome activation and the subsequent production of IL-1 β [41].

***Lycium barbarum* (LBP)**

It has been used as a traditional Chinese medicine to nourish liver, kidneys and the eyes. LBP attenuated ethanol-induced hepatic damages through suppressing the activation of NLRP3 inflammasome in a TXNIP-dependent manner in the BRL-3A in vitro system. These results further supported evidence that inhibition of hepatic TXNIP/NLRP3 inflammasome axis contributes to the alleviation of hepatic injury caused by ethanol [42].

Impatiens textori Miq.

I. textoria as a member of Balsaminaceae family, treated at 25, 50, and 100 µg/mL concentrations suppressed interleukin-1β secretion through the attenuation of NLRP3 inflammasome activation leading to the decreased amount of ASC oligomerization and caspase-1 maturation. It inhibited the NLRP3 expression and cell recruitment at the lung tissue in the ALI mouse model. Anti-inflammatory effects of *I. textoria* are via the attenuation of NLRP3 inflammasome activation [43].

Aloe vera

A. vera as an immunomodulatory agent inducing anti inflammatory effects. It mediated strong reduction of IL-1 appears to be the consequence of the reduced expression of both pro-IL-1 as well as NLRP3 inflammasome components via suppressing specific signal transduction pathways. The expression of the ATP sensor P2X7 receptor is also down regulated by Aloe vera that could also contribute to the attenuated IL-1 cytokine secretion. These results may provide a new therapeutic approach to suppression of inflammasome-mediated responses [44].

Syneilesis palmata (Thunb.)

S. palmate from Asteraceae family is a traditional Korean therapeutic herb widely used to treat pain, and arthritis. *S. palmate* inhibited the LPS-stimulated release of proinflammatory mediators, such as nitric oxide and interleukin IL-6 in RAW 264.7 cells. *S. palmate* treatment attenuated IL-1β secretion via the inhibition of NLRP3 inflammasome activation induced by monosodium urate, ATP, and nigericin. Further, *S. palmate* ameliorated the severity of NLRP3 inflammasome-mediated symptoms in LPS-induced endotoxin and E. coli-induced sepsis mouse models. Effects of *S. palmate* are mediated through the regulation of TRIF-dependent signaling and inflammasome activation [45].

Citral (3,7-dimethyl-2,6-octadienal)

Citral is a major active compound in a *Litsea cubeba*. Citral inhibited NLRP3 inflammasome activation and levels of ROS, NAD(P)H oxidase subunit p47phox, or COX-2, and it enhanced the activation of nuclear factor E2-related factor 2 (Nrf2). Citral alleviates severe lupus nephritis model via inhibition of the activation of NLRP3 inflammasome and enhanced activation of Nrf2 antioxidant signaling. Generation of reduced ROS by Citral may explain the resultant decreased pro-IL-1β protein levels [46].

Celastrol

Celastrol is a quinonemethide triterpene derived from a plant extract used in herbal medicine. It inhibited the proteasome-dependent degradation of proteins in RAW264.7 cells. It also blocked stimulation of IL-18 processing, indicating that celastrol acted upstream of inflammasome activation. Celastrol as an inhibitor of lethal toxin-mediated macrophage lysis and suggests an inhibitory mechanism involving inhibition of the proteasome pathway [47].

Schisandrin

Schisandrin is a phytochemical compound that can induce cellular antioxidant response. It also can suppress the inflammasome activation. Schisandrin B treatment (2 mmol/kg p.o.) ameliorated the Imject Alum-induced peritonitis, as indicated by suppressions of caspase1 activation and plasma IL-1β level [48].

Resveratrol

Resveratrol as a natural compound found in many types of plants and is one of the most intensely investigated phytochemicals. It can inhibit NLRP3-inflammasome activation via inhibiting the accumulation of acetylated α-tubulin. In addition, resveratrol can inhibit the acetylated-α-tubulin-mediated spatial arrangement of mitochondria and their subsequent contact with the endoplasmic reticulum. Actually, resveratrol targets assembly of NLRP3 and ASC, thus an increasing dose of resveratrol suppress mitochondrial ROS production, and subsequent NLRP3-inflammasome activation [49].

Dehydrodiconiferyl alcohol (DHCA)

DHCA is a lignan compound from *Cucurbita moschata*, reduced the production of pro-inflammatory cytokines such as TNF- α , IL-6, IL-1 β and CCL2 and mediators such as iNOS, COX-2 and ROS. DHCA suppressed activation of inflammasomes by down-regulating the activity of I- κ B kinase and DNA binding activity of NF- κ B [50].

Luteoloside

Luteoloside is a flavonoid isolated from some plants such as *Gentiana macrophylla*. Luteoloside can significantly decrease the intracellular reactive oxygen species accumulation. The decreased levels of ROS induced by luteoloside were accompanied by decrease in expression of NLRP3 inflammasome resulting in decrease in proteolytic cleavage of caspase-1. Inactivation of caspase-1 through luteoloside resulted in inhibition of IL-1 β . Thus, luteoloside exerts its inhibitory effect through inhibition of NLRP3 inflammasome [51].

Sulforaphane

Sulforaphane, as natural compound in cruciferous vegetables, increased nuclear factor erythroid 2 related factor 2 (Nrf2) nuclear translocation which results in decrease of IL-1 β release and microRNA-146a production. Also effects of sulforaphane were replicated by the tyrosine kinase inhibitor, herbimycin A, and Nrf2 activator. These findings suggest that sulforaphane has a potential application in suppression of inflammasome activation [52].

Pulsatilla decoction

Pulsatilla decoction as a Chinese formula is likely to exert their therapeutic effect by activating NLRP3 inflammasome which further promotes the formation of the corresponding inflammation factors such as IL-18 and IL-1 β [53].

Wuling San

Wuling San, a famous prescription in Chinese medicine, is composed of some natural compounds such as polyporus (a genus of fungi), poria (a famous home-grown bulk Chinese herb), *Alismatis rhizoma*, *Cinnamomi cortex* and *Atractylodis macrocephalae* rhizome. Wuling San suppressed the activation of TLR4/MyD88 signaling to inhibit NF- κ B signaling and mitogen-activated protein kinases activation in the kidney of fructose-fed mice. Additionally, Wuling San decreased NLRP3 inflammasome activation and IL-1 β secretion [54].

Regarding the significance of inflammatory diseases, and the necessity of preventing and treating these diseases, the aim of this review article is to report medicinal plants and their nature-based derivatives that are effective on inflammasome complex. Agents that reduce the production and activity of proinflammatory cytokines are likely to have clinical applications. For this purpose, future studies should examine which components in the extracts are responsible for the anti-inflammatory properties described in medicinal plants. Also, further studies are needed to better understand the mechanism by which medicinal plants inhibit inflammasome formation. The formation and activation of inflammasome complex is followed by the release of two cytokines, IL-1 β and IL-18, from cells [55]. Among the inflammasomes, the NLRP3 inflammasome is the most widely studied and is known to be activated by a wide range of PAMPs and DAMPs such as nucleotide ATP, nigericin, and monosodium urate. NLRP3 inflammasome, known to be activated upon cellular 'danger' signals, triggers innate immune defense through the secretion of pro-inflammatory cytokines, such as IL-1 β and IL-18 [3, 56]. Therefore, IL-1 β and inflammasomes have emerged as therapeutic targets for an expanding number of systemic and local inflammatory conditions. NLRP3 was recently identified to form a cytoplasmic complex known as the NLRP3 inflammasome, which potently modulates innate immune function by regulating the maturation and secretion of pro-inflammatory cytokines, such as IL-1 β [57]. IL-1 together with TNF are among the most powerful cytokines that under pathological conditions could cause cell degeneration and cell death resulting in multiple organ dysfunction [58]. Activation of the NLRP3 inflammasome is dependent on the generation of ROS [59, 60]. In fact, all known NLRP3 activators generate ROS and, conversely, inhibitors of ROS block inflammasome activation [61]. Our study demonstrated that some medicinal plants were found to

suppress NLRP3 inflammasome activation. This study was the first review article to show scientific evidence and justify the traditional use medicinal plants in the treatment of inflammation related disorders through block inflammasome activation. The strong protective effects exhibited by medicinal plants in both in vitro and in vivo experimental models might be partly mediated by the regulation of key inflammatory mechanisms, such as the TRIF signaling pathway and NLRP3 inflammasome activation, providing a valuable therapeutic or/and prophylactic strategy by controlling inflammation-mediated disorders

Table 1: The effects of medicinal plants and their derivatives on inflammasome

Plants or their derivatives	Study design and dosage	Results	Ref.
<i>Sophora flavescens</i>	<i>In vitro</i> and <i>In vivo</i> ; 50 ng/ μ	Blocked the phosphorylation of IKK α / β , and the cleavage of caspase-1, a key component of the inflammasome.	[41]
<i>Lycium barbarum</i>	<i>In vitro</i> ; 50 g/ml	Suppressed the activation of NLRP3 inflammasome in a TXNIP-dependent manner.	[42]
<i>Impatiens textori</i> Miq.	<i>In vivo</i> ; 25, 50, and 100 μ g/mL	Suppressed the interleukin-1 β secretion through the attenuation of NLRP3 inflammasome activation.	[43]
<i>Aloe vera</i>	<i>In vitro</i> ; 1, 3, and 10 v/v%	Inhibited the expression of pro-IL-1, Nlrp3, caspase-1 as well as that of the P2X7 receptor.	[44]
<i>Syneilesis palmata</i> (Thunb.) Maxim	<i>In vitro</i> and <i>In vivo</i> ; 100 μ g/mL	Regulated the TRIF-dependent signaling and inflammasome activation.	[45]
Citral	<i>In vitro</i> and <i>In vivo</i> ; 200 mg/kg	Inhibited the activation signal of NLRP3 and enhanced activation of Nrf2.	[46]
Celastrol	<i>In vitro</i> ; 10 mM	Blocked the stimulation of IL-18 processing.	[47]
Schisandrin	<i>In vitro</i> and <i>In vivo</i> ; 2 mmol/kg p.o	Suppressed the caspase1 activation and plasma IL-1 β level.	[48]
Resveratrol	<i>In vitro</i> ; low < 5 μ M, and higher > 25 μ M	Inhibited the acetylated- α -tubulin- causing insufficient assembly of ASC on the mitochondria and NLRP3 on the endoplasmic reticulum.	[49]
Luteoloside	<i>In vitro</i> and <i>In vivo</i> ; 25, and 50 mM	Reduced the intracellular reactive oxygen species (ROS) accumulation.	[51]
Sulforaphane	<i>In vitro</i> ; 5 mM	Dephosphorylated the STAT-1 and activated the Nrf2/HO-1 cascades.	[52]
Wuling San	<i>In vitro</i> and <i>In vivo</i> ; 987, 1316, 1755 and 2340 mg/kg	Suppressed the TLR4/MyD88 signaling and NLRP3 inflammasome activation to reduce IL-1 β production	[54]

Conclusion

Medicinal plants through their active ingredients have a good therapeutic effect of diseases and disorders (62-112). Medicinal plants and their derivatives down regulate pro-inflammatory cytokine production. Because patients with rheumatoid arthritis, septic shock, autoimmune disorders, tumors, Alzheimer's disease, and chronic inflammatory diseases secrete large amounts of IL-1 β , the agents that decline the production and activity of this proinflammatory cytokine can have clinical applications. Suppression of inflammasome might be an important therapeutic strategy to improve inflammatory-mediated diseases. Therefore, identifying the compounds that regulate the NLRP3-inflammasome is important to develop therapeutic methods with which infectious and inflammatory diseases are treated.

Conflict of Interests

There is no any conflict of interest.

References

1. Lamkanfi M, Dixit VM. Inflammasomes and their roles in health and disease. *Annu Rev Cell Develop Biol* 2012;28:137-61.
2. Sani MRM, Moghaddam MM, Aghamollaei H, Hassanpour K, Taheri RA, Farnoosh G. Investigation of caspase-1 activity and interleukin-1 β production in murine macrophage cell lines infected with *Leishmania major*. *Asian Pac J Trop Med* 2014;7:70-3.
3. Schroder K, Tschopp J. The inflammasomes. *Cell* 2010;140:821-32.
4. Fata A, Mahmoudian M, Varasteh A, Sankian M. Monarch-1 activation in murine macrophage cell line (J774 A. 1) infected with Iranian strain of *Leishmania major*. *Iran J Parasitol* 2013;8:207.
5. Sagulenko V, Thygesen SJ, Sester DP, Idris A, Cridland JA, Vajjhala PR, et al. AIM2 and NLRP3 inflammasomes activate both apoptotic and pyroptotic death pathways via ASC. *Cell Death Differ* 2013;20:1149-60.
6. Lamkanfi M, VandeWalle L, Kanneganti TD. Deregulated inflammasome signaling in disease. *ImmunolRev* 2011;243:163-73.
7. Guo H, Callaway JB, Ting JP. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med* 2015;21:677-87.
8. Sani MM, Hajizade A, Sankian M, Fata A, Mellat M, Hassanpour K, et al. Evaluation of the expression of the inflammasome pathway related components in *Leishmania major*-infected murine macrophages. *Eur J ExpBiol* 2013;3:104-9.
9. Hua KF, Yang SM, Kao TY, Chang JM, Chen HL, Tsai YJ, et al. Osthole mitigates progressive IgA nephropathy by inhibiting reactive oxygen species generation and NF-kappaB/NLRP3 pathway. *PloS one* 2013;8:e77794.
10. Lin J, Bi L, Yu X, Kawai T, Taubman MA, Shen B, et al. *Porphyromonasgingivalis* exacerbates ligature-induced, RANKL-dependent alveolar bone resorption via differential regulation of Toll-like receptor 2 (TLR2) and TLR4. *Infect Immun* 2014;82:4127-34.
11. Liao PC, Chao LK, Chou JC, Dong WC, Lin CN, Lin CY, et al. Lipopolysaccharide/adenosine triphosphate-mediated signal transduction in the regulation of NLRP3 protein expression and caspase-1-mediated interleukin-1 β secretion. *InflammRes* 2013;62:89-96.
12. Aruna R, Geetha A, Suguna P. Rutin modulates ASC expression in NLRP3 inflammasome: a study in alcohol and cerulein-induced rat model of pancreatitis. *MolCell Biochem* 2014;396:269-80.
13. Chen GY, Nunez G. Sterile inflammation: sensing and reacting to damage. *Nat Rev Immunol* 2010;10:826-37.
14. Masters SL, Simon A, Aksentijevich I, Kastner DL. Horror autoinflammaticus: the molecular pathophysiology of autoinflammatory disease. *AnnuRev Immunol* 2009;27:621.
15. Stienstra R, Joosten LA, Koenen T, van Tits B, van Diepen JA, van den Berg SA, et al. The inflammasome-mediated caspase-1 activation controls adipocyte differentiation and insulin sensitivity. *Cell Metabol* 2010;12:593-605.
16. Wen H, Gris D, Lei Y, Jha S, Zhang L, Huang MT-H, et al. Fatty acid-induced NLRP3-ASC inflammasome activation interferes with insulin signaling. *Nat Immunol* 2011;12:408-15.
17. Vandanmagsar B, Youm Y-H, Ravussin A, Galgani JE, Stadler K, Mynatt RL, et al. The NLRP3 inflammasome instigates obesity-induced inflammation and insulin resistance. *Nat Med* 2011;17:179-88.
18. Rabiei Z, Bigdeli M, Asadi M. The Effect of dietary virgin olive oil on brain lipid levels and brain edema in rat stroke models. *ZUMS Journal* 2013;21:56-64.
19. Kooti W, Ghasemiboroon M, Ahangarpour A, Hardani A, Amirzargar A, Asadi-Samani M. The effect of hydro-alcoholic extract of celery on male rats in fertility control and sex ratio of rat offspring. *J Babol Univ Med Sci* 2014; 16:43-9.
20. Khodadadi S, Rafeian-Kopaei M. Herbs, health and hazards; a nephrology viewpoint on current concepts and new trends. *Ann Res Antioxid*. 2016; 1:e05.
21. Shirani M, Alibabaei Z, Kheiri S, Shirzad H, Tajiri F, Asgari A, et al. Effect of *Euphorbia helioscopia* extract on acute and chronic pain in mice. *J Babol Univ Med Sci* 2011;13: 14-18.
22. Beyrami-Miavagi A, Farokhi F, Asadi-Samani M. A study of the effect of prostodin and hydroalcoholic extract of *Malvaneglecta* on kidney histopathology and renal factors in female rats. *Adv Environ Biol* 2014:942-8.

23. Bahmani M, Shirzad H, Rafieian S, Rafieian-Kopaei M. Silybummarianum: Beyond Hepatoprotection. *J Evid Based Complementary Altern Med* 2015;20: 292-301.
24. Samarghandian S, Asadi-Samani M, Farkhondeh T, Bahmani M. Assessment the effect of saffron ethanolic extract (*Crocus sativus* L.) on oxidative damages in aged male rat liver. *Der Pharm Lett* 2016;8:283-90.
25. Mahmoudian Sani M, Asadi-Samani M, Rouhi-Boroujeni H, Banitalebi-Dehkordi M. Phytopharmacology and phytotherapy of regulatory T cells: A new approach to treat multiple sclerosis. *Der Pharm Lett* 2016;8:215-20.
26. Parsaei P, Bahmani M, Karimi M, Naghdi N, Asadi-Samani M, Rafieian-Kopaei M. A review of analgesic medicinal plants in Iran. *Der Pharm Lett* 2016;8:43-51.
27. Jivad N, Bahmani M, Asadi-Samani M. A review of the most important medicinal plants effective on wound healing on ethnobotany evidence of Iran. *Der Pharm Lett* 2016;8:353-7.
28. Akhlaghi M, Shabanian G, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. Citrus aurantium Blossom and Preoperative Anxiety. *Rev Bras Anesthesiol* 2011;61:702-12.
29. Parsaei P, Bahmani M, Naghdi N, Asadi-Samani M, Rafieian-Kopaei M, Boroujeni S. Shigellosis phytotherapy: A review of the most important native medicinal plants in Iran effective on Shigella. *Der Pharm Lett* 2016;8:249-55.
30. Jivad N, Asadi-Samani M, Moradi M. The most important medicinal plants effective on migraine: A review of ethnobotanical studies in Iran. *Der Pharm Chem* 2016;8:462-6.
31. Nasri H, Sahinfard N, Rafieian M, Rafieian S, Shirzad M, Rafieian-kopaei M. Turmeric: A spice with multifunctional medicinal properties. *J HerbmedPharmacol* 2014; 3: 5-8.
32. Mohsenzadeh A, Ahmadipour S, Ahmadipour S, Asadi-Samani M. Iran's medicinal plants effective on fever in children: A review. *Der Pharm Lett* 2016;8:129-34.
33. Ahmadipour S, Ahmadipour S, Mohsenzadeh A, Asadi-Samani M. The importance of some native medicinal plants of Iran effective on gastrointestinal disorders in children: A review. *Der Pharm Lett* 2016;8:61-6.
34. Kafeshani M. Ginger, micro-inflammation and kidney disease. *J Renal Endocrinol*.2015; 1:e04.
35. Rafieian-Kopaei M, Nasri H. Herbal antioxidant therapy in dialysis patients. *Ann Res Dial* 2016; 1:e02.
36. Baradaran A, Nasri H, Rafieian-Kopaei M. Comment on: Anti-oxidative stress activity of stachyslavandulifolia aqueous extract in humans. *Cell J* 2013;15:272-3.
37. Madihi Y, Merrikhi A, Baradaran A, Rafieian-Kopaei M, Shahinfard N, Ansari R, et al. Impact of Sumac on postprandial high-fat oxidative stress. *Pak J Med Sci* 2013; 29: 340-5.
38. Asadi-Samani M, Kooti W, Aslani E, Shirzad H. A Systematic Review of Iran's Medicinal Plants With Anticancer Effects. *J Evid Based Complementary Altern Med* 2016;21:143-53.
39. Parsaei P, Bahmani M, Naghdi N, Asadi-Samani M, Rafieian-Kopaei M, Tajeddini P, et al. Identification of medicinal plants effective on common cold: An ethnobotanical study of Shiraz, South Iran. *Der Pharm Lett* 2016;8:90-7.
40. Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Ethanolic extract of garlic for attenuation of gentamicin-induced nephrotoxicity in wistar rats. *Iran J Kidney Dis* 2013;7:376-82.
41. Lee JH, Shin H, Kim YJ, Paek SH, Jin S, Ha UH. Pseudomonas aeruginosa-induced IL-1beta production is inhibited by Sophoraflavescens via the NF-kappaB/inflammasome pathways. *J Microbiol* 2014;52:1044-9.
42. Xiao J, Zhu Y, Liu Y, Tipoe GL, Xing F, So KF. Lyciumbarbarum polysaccharide attenuates alcoholic cellular injury through TXNIP-NLRP3 inflammasome pathway. *Int J BiolMacromol* 2014;69:73-8.
43. Sun X, Shim DW, Han JW, Shin WY, Sim EJ, Kim MK, et al. Anti-inflammatory effect of Impatiens textoriMiq. extract via inhibition of NLRP3 inflammasome activation in in vitro and in vivo experimental models. *J Ethnopharmacol* 2015;170:81-7.
44. Budai MM, Varga A, Milesz S, Tozser J, Benko S. Aloe vera downregulates LPS-induced inflammatory cytokine production and expression of NLRP3 inflammasome in human macrophages. *MolImmunol* 2013;56:471-9.
45. Han JW, Shim DW, Shim EJ, Kim MK, Shin YK, Kwak SB, et al. Syneilesispalmata (Thunb.) Maxim. extract attenuates inflammatory responses via the regulation of TRIF-dependent signaling and inflammasome activation. *J Ethnopharmacol* 2015;166:1-4.
46. Kassem A, Henning P, Lundberg P, Souza PP, Lindholm C, Lerner UH. Porphyromonasgingivalis Stimulates Bone Resorption by Enhancing RANKL (Receptor Activator of NF-kappaB Ligand) through Activation of Toll-like Receptor 2 in Osteoblasts. *J BiolChem* 2015; 290:20147-58.

47. Chapelsky S, Batty S, Frost M, Mogridge J. Inhibition of anthrax lethal toxin-induced cytolysis of RAW264.7 cells by celastrol. *PloS one* 2008;3:e1421.
48. Leong PK, Ko KM. Schisandrin B induces an Nrf2-mediated thioredoxin expression and suppresses the activation of inflammasome in vitro and in vivo. *BioFactors* (Oxford, England) 2015;41:314-23.
49. Misawa T, Saitoh T, Kozaki T, Park S, Takahama M, Akira S. Resveratrol inhibits the acetylated alpha-tubulin-mediated assembly of the NLRP3-inflammasome. *IntImmunol*2015;27:425-34.
50. Lee J, Choi J, Kim S. Effective suppression of pro-inflammatory molecules by DHCA via IKK-NF-kappaB pathway, in vitro and in vivo. *Br J Pharmacol*2015;172:3353-69.
51. Fan SH, Wang YY, Lu J, Zheng YL, Wu DM, Li MQ, et al. Luteoloside suppresses proliferation and metastasis of hepatocellular carcinoma cells by inhibition of NLRP3 inflammasome. *PloS one* 2014;9:e89961.
52. An YW, Jhang KA, Woo SY, Kang JL, Chong YH. Sulforaphane exerts its anti-inflammatory effect against amyloid-beta peptide via STAT-1 dephosphorylation and activation of Nrf2/HO-1 cascade in human THP-1 macrophages. *NeurobiolAging* 2016;38:1-10.
53. Wang ZL, Sun PP, Li TX, Li SX. Pulsatilladecocotion treat inflammatory bowel disease by activating NLRP3. *Zhong Yao Cai*2012;35:1280-6.
54. Yang Y, Zhang DM, Liu JH, Hu LS, Xue QC, Ding XQ, et al. Wuling San protects kidney dysfunction by inhibiting renal TLR4/MyD88 signaling and NLRP3 inflammasome activation in high fructose-induced hyperuricemic mice. *J Ethnopharmacol*2015;169:49-59.
55. Keyel PA. How is inflammation initiated? Individual influences of IL-1, IL-18 and HMGB1. *Cytokine* 2014;69:136-45.
56. Nakaya M, Hara H, Yamazaki M, Yoshida H, Sankian M, Sani MM, et al., editors. Immunity to protozoan parasite infection (PP-002). *International Immunology Meeting Abstracts*; 2010: Oxford University Press.
57. Tschopp J, Schroder K. NLRP3 inflammasome activation: The convergence of multiple signalling pathways on ROS production? *Nat Rev Immunol* 2010;10:210-5.
58. Tracey KJ, Fong Y, Hesse DG, Manogue KR, Lee AT, Kuo GC, et al. Anti-cachectin/TNF monoclonal antibodies prevent septic shock during lethal bacteraemia. *Nature* 1987;330:662-4.
59. Sorbara MT, Girardin SE. Mitochondrial ROS fuel the inflammasome. *Cell Res* 2011;21:558-60.
60. Wen H, Gris D, Lei Y, Jha S, Zhang L, Huang MT, et al. Fatty acid-induced NLRP3-ASC inflammasome activation interferes with insulin signaling. *Nat Immunol*2011;12:408-15.
61. Zhou R, Tardivel A, Thorens B, Choi I, Tschopp J. Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat Immunol* 2010;11:136-40.
62. Nikfarjam M, Bahmani M, Naimi A. Native medicinal plants of Iran effective on Memory and Learning: A Review. *International Journal of PharmTech Research*, 2016; 9(5), 466-473.
63. Asadi-Samani M, Moradi MT, Bahmani M, Shahrani M. Antiviral medicinal plants of Iran: A Review of Ethnobotanical evidence. *International Journal of PharmTech Research*, 2016; 9(5), 427-434.
64. Moradi MT, Asadi-Samani M, Bahmani M. Hypotensive medicinal plants according to Ethnobotanical evidence of Iran: A Systematic Review. *International Journal of PharmTech Research* 2016; 9(5), 416-426.
65. Moradi MT, Asadi-Samani M, Bahmani M, Shahrani M. Medicinal plants used for liver disorders based on the Ethnobotanical documents of Iran: A Review. *International Journal of PharmTech Research* 2016; 9(5), 407-415.
66. Zailanie K, Kartikaningsih H, Kalsum U, Sanjaya YA. Fucoxanthin effects of pure sargassum filipendula extract toward HeLa cell damage. *International Journal of PharmTech Research*, 2015; 8(3), 402-407.
67. Rosidah, Poppy Anjelisa Z. Hasibuan. Cytotoxic effect of n-hexane, ethylacetate and ethanol extracts of *Plectranthus amboinicus*, (Lour.) Spreng. on HeLa and Vero cells lines. *International Journal of PharmTech Research*, 2014; 6(6), 1806-1809.
68. SyafruddinIlyas. Effect of methanolic *Momordica charantia* seed extract and depot medroxyprogesterone acetate (DMPA) to quantity and quality of rat sperm. *International Journal of PharmTech Research* 2014; 6(6), 1817-1823.
69. Janani J, Estherlydia D. Antimicrobial activities of punicagranatum extracts against oral microorganisms. *International Journal of PharmTech Research*, 2013; 5(3), 973-977.
70. Mallick M, Bose A, Mukhi S. Comparative evaluation of the antioxidant activity of some commonly used spices. *International Journal of PharmTech Research*, 2016, 9(1), 1-8.

71. Saravanan D, Radhakrishnan M. Antimicrobial activity of mangrove leaves against drug resistant pathogens. International Journal of PharmTech Research, 2016, 9(1), 141-146.
72. Anarthe SJ, Pravalika A, Malavika E, Ganga Raju M. Assessment of immunomodulatory activity of *Ficus benghalensis* Linn. aerial roots. International Journal of PharmTech Research, 2016; 9(1), 153-163.
73. Jaffat HS, Nasir AS. Effect of turmeric extract (*Curcuma longa*) on physiological parameters and neurotransmitters in rats treated by lithium carbonate. International Journal of PharmTech Research, 2016, 9(2), 89-97.
74. Ezema BE, Odoemelam EI, Agbo MO. Phytochemical and Antibiotic Evaluation of the Methanol Extract of *Loranthus micranthus* Linn Parasitic on *Kola Accuminata*. International Journal of PharmTech Research, 2016; 9(2), 176-181.
75. Daud N, Rosidah M, Nasution P. Antidiabetic activity of *Ipomoea batatas* L. leaves extract in streptozotocin-induced diabetic mice. International Journal of PharmTech Research, 2016, 9(3), 167-170.
76. Tambe VD, Bhambhar RS. Studies on diuretics and laxative activity of the *Hibiscus tiliaceus* Linn. bark extracts. International Journal of PharmTech Research, 2016; 9(3), 305-310.
77. Jain PK, Das D, Jain P. Evaluating hair growth activity of herbal hair oil. International Journal of PharmTech Research, 2016, 9(3), 321-327.
78. Bhangale JO, Acharya NS, Acharya SR. Neuroprotective effect of pet ether extract of *Ficus religiosa* (L.) leaves in 3-nitropropionic acid induced Huntington disease. International Journal of PharmTech Research, 2015; 8(10), 57-69.
79. Shaikh AM, Shrivastava B, Apte KG, Navale SD. Effect of aqueous extract of *Curcuma zedoaria* and *Gloriosa superba* against DMH-Induced colon carcinogenesis in Wistar rats. International Journal of PharmTech Research, 2015; 8(10), 88-94.
80. Habeeb RA, Majed N. Some herbal medicinal plants activity against *Candida* spp which resistance to antifungal drugs. International Journal of PharmTech Research, 2015; 8(10), 146-150.
81. Al-Terehi M, Al-Saadi AH, Zaidan HK, Behjet RH, Haleem Z. Some plants extracts Synergism effects in Pathogenic bacteria, International Journal of PharmTech Research, 2015, 8(10), 158-164.
82. Ragavan Balliah, Monisha Sudhakar(2015), In Vitro evaluation of cytotoxic and antiproliferative activity of a polyherbal extract against H9c2 cardiac cells. International Journal of PharmTech Research, 2015; 8(10), 191-197.
83. Jasmine R, Sakthivel H. Role of compounds from *Terminalia chebula* exhibiting anti-cholesterol property. International Journal of PharmTech Research, 2015; 8(10), 210-215.
84. Jansen Silalahi, Dwi Pertiwi, Aminah Dalimunthe, Yosy C E Silalahi(2015), Effect of acute consumption of coconut and palm oil on swimming capacity endurance of mice (*Mus musculus*). International Journal of PharmTech Research, 2015, 8(9), 55-59.
85. Kamali H, Ahmadzadeh Sani T, Feyzi P, Mohammadi A. Chemical composition and antioxidant activity from essential oil of *Capsella bursa-pastoris*. International Journal of PharmTech Research, 2015; 8(8), 1-4.
86. Velu V, Malipeddi H. In vitro anti-arthritis and hemolytic activity of leaf extracts of *Tragia involucrata*. International Journal of PharmTech Research, 2015; 8(7), 46-50.
87. Himaja M, Das P. Antioxidant, anti-arthritis and hypoglycemic activity of *Oxalis corniculata* Linn. leaf extracts. International Journal of PharmTech Research, 2015; 8(7), 51-57.
88. Duraisankar M, Devi M, Shanmugasundram P. Hepatoprotective activity of alcoholic extract of *Chonemorpha fragrans* root in against Paracetamol and Isoniazid-induced liver damage in rats. International Journal of PharmTech Research, 2015; 8(7), 232-242.
