

## **Synthesis, Structural Characterization, and Catalytic Properties of Cu(II), Ru(II) and Pd(II) Complexes with Nitrogen & Oxygen donor Tetradentate Schiff base ligands**

**P. Venugopal<sup>1</sup>, S. Bhaskar<sup>3</sup>, and P. Muralidhar Reddy<sup>2,3\*</sup>**

<sup>1</sup>Department of Chemistry, Vasavi College of Engineering, Ibrahimbagh, Hyderabad

<sup>2</sup>Toxicology and Biomedicine Research Group, Faculty of Applied Sciences, Ton Duc Thang University, Ho Chi Minh City, Vietnam

<sup>3</sup>Department of Chemistry, Nizam College, Osmania University, Hyderabad, India

**Abstract** : Cu (II), Pd (II) and Ru (II) metal complexes of N<sub>4</sub> and N<sub>2</sub>O<sub>2</sub> based tetradentate Schiff base ligands were prepared via non-template method by the condensation of 3, 5-diaminobenzoic acid with o-salicylaldehyde (HL) and o-phthalaldehyde (OPA) (HL<sup>1</sup>) respectively. The synthesized ligands and metal complexes were well characterized by a variety of techniques such as elemental analysis, FTIR, <sup>1</sup>H NMR, MS, electronic spectroscopy, and conductance measurements. The catalytic activity of Schiff base metal complexes were screened in the reduction of dabigatran, which is an intermediate in the morphine synthesis, and C-N cross coupling reactions resulting nilutamide is an anticancer drug, abacavir that is antiviral drug. The Schiff base metal complexes can catalyze selectively and shows moderate to high yields.

**Key words** : Schiff bases ligands, metal complexes, characterization, catalysis.

### **Introduction**

Schiff bases were versatile compounds synthesized from the condensation of a variety of readily available aldehydes and primary amines and regarded as privileged ligands due to their capability to form complexes with different transition metals<sup>1-26</sup>. Schiff base metal complexes have wide range of applications in many fields including analytical<sup>27, 28</sup>, medicinal<sup>29, 30</sup>, organic<sup>31</sup>, inorganic chemistry. These complexes can act as catalysts in several reactions such as polymerization<sup>32</sup>, oxidation<sup>33, 34</sup>, hydrolysis<sup>14, 35</sup> reduction<sup>36</sup>, epoxidation<sup>37</sup>, hydrosilylation<sup>38</sup> Henry reaction<sup>39</sup>, C-N<sup>40</sup>, C-S, Kumada Corriu cross coupling<sup>41</sup> reactions, and C-H activation of various substrates. Schiff base metal complexes have great importance in industrial applications and in various biological processes including photosynthesis, dioxygen transport, and potential applications as metal extractants, radiotherapeutic, medical imaging agents and anti-tumour therapeutic agents<sup>42</sup>. For the past few years, several investigations have been done on the coordination chemistry of Schiff base coordinated transition metal complexes and supramolecules in order to mimic the physical and chemical properties and applied in biological processes. Macrocyclic Schiff base metal complexes derived from o-phthalaldehyde (OPA) were recently reported by Shakir and co-workers via template method<sup>43</sup>. To the best of our knowledge, there are very few reports on the synthesis and catalytic applications of Schiff base ligated transition metal (II) compounds derived from o-phthalaldehyde. In the present work, in continuation to earlier studies towards the synthesis of novel coordination compounds and their catalytic activities, we have synthesized N<sub>4</sub> and N<sub>2</sub>O<sub>2</sub> tetradentate ligands from o-phthalaldehyde by non-template method, and their complexes with copper, palladium

and ruthenium. Catalytic activities of metal complexes were investigated in reduction of dabigatran, which is an intermediate in the synthesis of morphine and C-N cross coupling reactions resulting nilutamide and abacavir. Dabigatran is new oral anticoagulant which inhibit factor thrombin and an alternative to warfarin in patients requiring cardioversion<sup>44</sup>. Nilutamide is a non-steroidal prostate anticancer drug and pneumotoxic, hepatotoxic nitroaromatic antiandrogen and used in the treatment of prostate carcinoma in man<sup>45</sup>. Abacavir is a nucleoside analogue reverse-transcriptase inhibitor of HIV type 1 replication and deoxy-guanosine base. Abacavir shows similar in vitro potency to other nucleoside analogues and an antiretroviral drug. Abacavir is associated with hypersensitivity in patients<sup>46</sup>.

## Material and measurements

### Materials


All the solvents, amines, aldehydes, and metal salts used in this study were of analar grade. The solvents were distilled by standard procedures before use. The ligands used in this work were prepared and characterized by IR, UV-Visible and <sup>1</sup>H-NMR spectral data. The water used in the study was essentially double distilled water. The solvents were distilled and stored over molecular sieves. The purity of the compounds was checked by TLC using Merck 60F254 silica gel plates

### Physical Measurements

The MPs of all the ligands were determined on a Buchi- 510 melting point apparatus. The MPs of the complexes could not be determined as the complexes were found to decompose before melting. UV-Visible spectra were recorded with Shimadzu UV-2140A, a UV-Visible double beam spectrophotometer with matched quartz cells of path length 1 cm at Osmania University, Hyderabad. The IR spectra were recorded in KBr pellets on Perkin Elmer-283 spectrophotometer at Osmania University, Hyderabad. The scanning rate was 6 min in the range of 4000-200 cm<sup>-1</sup>. <sup>1</sup>H NMR spectra were recorded in CDCl<sub>3</sub> or DMSO-d<sub>6</sub> on a Bruker AVAVCE III 400 MHz -NMR spectrometer using TMS as the internal standard at ICT Hyderabad. NMR spectral data provide valuable information regarding the structure of Schiff bases. The NMR spectra of the Metal (II) complexes were also recorded to know the nature of binding of metal ion.

### Synthesis of HBAB Schiff base ligand:

2 ml of salicylaldehyde dissolved in minimum quantity of methanol and added drop by drop to methanolic solution of 3,5 diamino benzoic acid (1.52 grams, 0.1moles) with continuous stirring and refluxed on water bath for about 1hour. Yellow colored precipitate was separated out. It was filtered off, washed with hot methanol and glacial acetic acid, and dried.



**HBAB(S):**C<sub>21</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>; ESI-MS m/z [M+1] 361; IR: HBAB(S): 3070, 1719, 1687, 1620, 1569, 1305, 751; <sup>1</sup>H-NMR: 6.97-7.02(3H), 7.41-7.46(5H), 7.69-7.72(3H, t), 7.76-7.77(3H, t), 7.84-7.86(2H, d).

### Preparation of metal complexes

#### Preparation of Cu (II) complex with HBAB:

0.0025 moles HBAB dissolved in 50:50 of chloroform and methanol. To this 0.0025 moles methanolic solution of copper (II) chloride added drop wise with continuous stirring on magnetic stirrer for about 30 minutes. A pista green coloured precipitate was separated out. It was filtered off, washed with hot methanol and 50:50 chloroform and methanol solution, and dried.

**Preparation of Pd (II) complex with HBAB:**


0.0025m moles HBAB dissolved in 50:50 of chloroform and methanol. To this hot solution, add 0.0025m moles of methanolic solution of Palladium (II) chloride with continuous stirring on magnetic stirrer for about 1 hour. Orange red, Black coloured precipitate respectively was separated out. It was filtered off, washed with hot methanol and 50:50 of chloroform and methanol, and dried.

**Preparation of Ru (II) complex With HBAB:**

0.0025m moles HBAB dissolved in 50:50 of chloroform and methanol. 0.0025m moles Ruthenium tri chloride hydrate was dissolved in methanol. This solution was added to hot solution of HBAB with continuous stirring on magnetic stirrer. Black colored precipitate was separated off which was filtered and dried.

**Synthesis of PMAA Schiff base ligand:**

The methanolic solution of ortho phthalaldehyde (1.34 gram 0.1 moles) is mixed with methanolic solution of three, 5-diamino benzoic acid (1.52 grams 0.1 moles) with continuous stirring. A yellow coloured solution was refluxed for about 2 hours and cooled to room temperature, which yielded a yellow coloured precipitate. It was filtered off, washed with hot methanol, and dried.



**PMAA(A):** C<sub>22</sub>H<sub>18</sub>N<sub>4</sub>O<sub>4</sub>; FTIR: 3360, 3221, 3076, 2919, 1685, 1635, 1384, 776, 726. <sup>1</sup>H-NMR: 4.96-5.07(3H, q), 6.33-6.40(1H, q), 6.59-6.67(1H, q), 6.92-6.94(1H, d), 7.16(2H, s), 7.46-7.77(6H, m), 8.51(1H, s). ESI-MS m/z [M+1] 403.

**Preparation of Cu (II) complex with PMAA:**

0.0025 moles of Cu<sup>2+</sup> was dissolved in methanol and added to hot DMSO solution of 0.0025 moles PMAA with continuous stirring on magnetic stirrer. Bluish green colored was formed and refluxed on water bath about ½ hour. Bluish green precipitate was separated off this was filtered and dried.

**Preparation of Pd (II) complex with PMMA:**

0.0025 moles of palladium (II) chloride was dissolved in chloroform and added drop wise in to hot DMSO solution of 0.0025 moles PMMA with continuous stirring on water bath. After 1 hour yellow colored precipitate was separated off this was filtered and dried.


**Preparation of Ru (II) complex with PMMA:**

0.0025 moles of PMMA were dissolved in DMSO. To this methanolic solution of 0.0025 moles ruthenium tri chloride was added drop wise with continuous stirring on hot plate. A black colored compound separated. This was filtered and dried.

## Catalysis

### Reduction of dabigatrin using Pd-Schiff's base Metal Complexes

#### Scheme:1




Dissolve Ethyl 3-(4-(methylamino)-3-nitro-N-(pyridin-2-yl) benzamido) propanoate (2.5gm) in ethyl acetate (25ml) in an autoclave. Add 250 mg of Pd/C at 25°C to 35°C and Pd Catalyst (Pd-HBAB & Pd-PMAA complex, 50mg) stirred for 20 min. Heat the reaction mixture at 50- 55°C under 5 Kg hydrogen pressure for five hrs. The reaction mixture was filtered and washed with ethyl acetate and distil the filtrate to get residue. Add 30 mL isopropanol to the residue and heat for 5-10 minutes at 70-75°C. The reaction mixture was cooled to 0-5°C and filtered. The wet-cake is washed with isopropanol and dried to afford 850 mg ethyl 3-(3-amino-4-(methylamino)-N-(pyridin-2-yl) benzamido) propanoate as a solid. Yield%-58 to 65

Mol Formula: C<sub>18</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>; IR: 3271, 3160, 2925, 2210, 1753, 1728, 1637, 1182, 1023, 894, 826. <sup>1</sup>H-NMR(DMSO-d<sub>6</sub>, 300 Hz's):1.09(3H, t), 2.65-2.70(2H, t), 3.75(3H, s), 3.93-4.0(2H, Q), 4.19-4.24(2H, t), 4.24-4.61(2H, d), 7.09-7.17(2H, m), 7.26-7.29(1H, t), 7.39-7.57(5H, m); ESI-MS m/z [M+1] 343.

### C-N cross coupling reactions using Cu-Schiff's base Metal Complexes

#### Scheme-2



#### Procedure with Copper Catalyst-PMAA complex

Dissolve two-(trifluoromethyl)-4-iodo-1-nitrobenzene (2gm) in N, N-dimethylformamide (5ml) at 25-30°C, add 5, 5-dimethylimidazolidine-2, 4-dione (1.3gm) and CuO (250mg), Copper catalyst-1 (40mg) in to the mixture. Heat the mixture to 100-110°C and maintained at 100-110°C until two-(trifluoromethyl)-4-iodo-1-nitrobenzene should be absent completely by TLC (30% EtOAc in Hexane), it takes 10hrs for completion of the reaction. Filter the reaction mass over hyflo bed and washed with DMF (1ml), quench the filtrate in to the water and stir for 30mints to get precipitate. Filter the solid and wash with water. Dissolve the solid in tert-butyl methyl ether, wash the organic layer with 10% EDTA solute on, and dry the organic layer over sodium sulfate. Distill the solvent up to ¼ of volume and crystallize the product at 0-5°C. Filter the solid and dry at 60-65°C. Wt. of the product: 1.2gm


Mol Formula: C<sub>12</sub>H<sub>10</sub>F<sub>3</sub>N<sub>3</sub>O<sub>4</sub>; IR: 3283, 2988, 1778, 1722, 1540, 1318, 1138. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 300 Hz's):1.44(6H, s), 8.07-8.09(1H, Q), 8.20-8.21(1H, d), 8.30-8.32(1H, d), 8.84(1H, s); ESI-MS m/z [M+1] 317.

#### Procedure with Copper Catalyst- HBAB complex

Dissolve 2-(trifluoromethyl)-4-iodo-1-nitrobenzene (1gm) in N, N-dimethylformamide (5ml) at 25-30°C, add 5, 5-dimethylimidazolidine-2, 4-dione (0.65gm) and CuO (125mg), Copper catalyst-1 (20mg) in to the mixture. Heat the mixture to 100-110°C and maintained at 100-110°C till 2-(trifluoromethyl)-4-iodo-1-nitrobenzene should be absent completely by TLC (30% EtOAc in Hexane), it takes 10hrs for completion of the

reaction. Filter the reaction mass over hyflo bed and washed with DMF (1ml), quench the filtrate in to the water and stir for 30mints to get precipitate. Filter the the solid and wash with water. Dissolve the solid in tert-butyl methyl ether and wash the organic layer with 10% EDTA solute on and dry the organic layer over sodium sulfate. Distill the solvent up to ¼ of volume and crystallize the product at 0-5°C. Filter the solid and dry at 60-65°C. Wt. of the product: 0.55gm

### Scheme-3



### Procedure:

Add ((1S, 3R)-4-(2-amino-6-chloro-9H-purin-9-yl) cyclopent-2-enyl) methanol (2gm), NaHCO<sub>3</sub> (475 mg, 5.65 mmol), cyclopropylamine (2 ml, 28.3 mmol) and Copper Catalyst (Cu-HBAB & Cu-PMAA complex, 45mg) in to Isopropanol (15ml). Stir the mixture for 30-35mints at 80-82°C. Filter the salts. Distill off the filtrate, isopropanol (10 ml) was added and concentrated twice to dryness. The syrup was dissolved in hot isopropanol (15 ml). This mixture is cooled to 0-2<sup>0</sup> C and filters the resulting slurry. Dry the solid under vacuum at 30-35°C. Wt. of compound: 1.25-1.3gm

Mol Formula: C<sub>14</sub>H<sub>18</sub>N<sub>6</sub>O IR: <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 300 Hz's):0.6(2H, ), 0.7(2H, ), 1.55(1H, ), 2.0(1H, ), 2.6-2.7(1H, ), 2.8(1H, ), 3.1(1H, ), 3.40(1H, ), 3.45(2H, ), 4.10(1H, ), 5.4(1H, ), 5.85(3H, ), 6.15(1H, ), 7.3(1H, ), 7.7(1H, ); ESI-MS m/z [M+1] 287.

### Results and discussion

In the present investigations, six new Schiff base Cu (II), Ru (II), Pd (II), complexes were synthesized by treating Metal chlorides with two Schiff base ligands separately using non-template method and all the complexes are stable in air and soluble in various organic solvents. The purity of the Schiff base ligands were monitored by TLC. The percentages of carbon, hydrogen and nitrogen were determined experimentally using CHN analyzer. The physical and analytical data for the newly synthesized compounds is in good agreement with the proposed molecular formulae (Table 1)

**Table 1 Analytical data of Schiff base ligands and their Cu (II), Ru (II), and Pd (II) metal complexes**

| Schiff base complex           | $\Lambda M^a$<br>( $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$ ) | Found (Cald (%)) |             |               |               |
|-------------------------------|--|------------------|-------------|---------------|---------------|
|                               |  | C                | H           | N             | M             |
| 1. HBAB                       | -  | 69.95 (69.99)    | 4.49 (4.48) | 7.78 (7.77)   | -             |
| 2. PMAA                       | -  | 65.67 (65.66)    | 4.50 (4.51) | 13.90 (13.92) | -             |
| 3. [Cu(HBAB)Cl <sub>2</sub> ] | 14.0   | 50.98 (50.97)    | 3.27 (3.26) | 5.67 (5.66)   | 12.82 (12.84) |
| 4. [Cu(PMAA)Cl <sub>2</sub> ] | 15.2   | 49.23 (49.22)    | 3.39 (3.38) | 10.45 (10.44) | 11.83 (11.84) |
| 5. [Ru(HBAB)Cl <sub>2</sub> ] | 16.0   | 47.39 (47.38)    | 3.05 (3.03) | 5.27 (5.26)   | 18.98 (18.99) |
| 6. [Ru(PMAA)Cl <sub>2</sub> ] | 15.5   | 46.03 (46.00)    | 3.16 (3.16) | 9.74 (9.75)   | 17.61 (17.60) |
| 7. [Pd(HBAB)] Cl <sub>2</sub> | 59.0   | 46.90 (46.91)    | 3.01 (3.00) | 5.20 (5.21)   | 19.78 (19.79) |
| 8. [Pd(PMAA)] Cl <sub>2</sub> | 63.0   | 45.59 (45.58)    | 3.14 (3.13) | 9.67 (9.66)   | 18.37 (18.36) |

### Infrared spectral analysis

In the IR spectra of the both ligands, a medium intensity band in the range of 1630-1660  $\text{cm}^{-1}$  is assigned to azomethine (C=N) and disappearance of the bands characteristic of the (-NH<sub>2</sub>) amine group at 3250  $\text{cm}^{-1}$  of 3,5-diaminobenzoic acid and carbonyl group (-C=O) at 1680  $\text{cm}^{-1}$  of o-phthalaldehyde suggests the formation of Schiff bases. Bands appearing in the region of 1492-1455, 1144-1033 and 799-751  $\text{cm}^{-1}$  were assigned to aromatic ring vibrations. A broad medium intensity band at 3274-3349 in PMMA ligand ascribed to (O-H) stretching frequency of aromatic carboxylic acid. The coordination of azomethine nitrogen was suggested by the negative shift of 20-50  $\text{cm}^{-1}$  of (C=N) stretch. This was further supported by a medium intensity band in the region of 430-450  $\text{cm}^{-1}$  that is ascribed to metal to nitrogen (M-N) stretching vibration. A band in the region of 530-570  $\text{cm}^{-1}$  assigned to (M-O) stretching vibration and band around 340  $\text{cm}^{-1}$  is due to (M-Cl) vibrational frequency. The selected spectral bands of ligands and metal complexes were presented in **Table.2 Important infrared frequencies**


| S No | Compound                   | (C=N) | (M-N) | (M-O) | (M-Cl) |
|------|----------------------------|-------|-------|-------|--------|
| 1.   | HL(HBAB)                   | 1620  | -     | -     | -      |
| 2.   | HL <sup>1</sup> (PMMA)     | 1635  | -     | -     | -      |
| 3.   | [Cu(HBAB)Cl <sub>2</sub> ] | 1590  | 430   | 550   | 320    |
| 4.   | [Cu(PMAA)Cl <sub>2</sub> ] | 1605  | 432   | -     | 340    |
| 5.   | [Ru(HBAB)Cl <sub>2</sub> ] | 1580  | 440   | 530   | 338    |
| 6.   | [Ru(PMAA)Cl <sub>2</sub> ] | 1610  | 445   | -     | 345    |
| 7.   | [Pd(HBAB)] Cl <sub>2</sub> | 1599  | 450   | 570   | -      |
| 8.   | [Pd(PMAA)] Cl <sub>2</sub> | 1590  | 448   | -     | -      |

### NMR spectral analysis

In <sup>1</sup>H-NMR spectra of the ligand PMMA a signal due to four equivalent CH=N [4H] protons appeared at 8.5 ppm. A broad peak in the range of 4.96-5.07 ppm is assigned to four equivalent aromatic (C-NH) protons and all the peaks in the range of 6.33-7.77 ppm are corresponding to [10H] aromatic protons. <sup>1</sup>H-NMR spectra of N<sub>2</sub>O<sub>2</sub> ligand (HBAB) shows a signal at 7.84-7.86 is due to azomethyne (CH=N) protons [2H] appeared in the range 7.84-7.87 ppm suggests the formation of Schiff base ligands. Additionally, in the spectra of Ru (II), and Pd (II) complexes the signal was moved downfield, this suggests the coordination of imino nitrogen to metal ion. No appreciable change is seen in the peak positions corresponding to aromatic protons and NH protons and hydroxy protons in the complexes.

### Electronic spectral analysis

The electronic spectra of Cu (II) complexes reported here show two characteristic bands in the range 660–870 (<sup>2</sup>B<sub>1g</sub> → <sup>2</sup>A<sub>1g</sub>) and 430–515 nm (<sup>2</sup>B<sub>1g</sub> → <sup>2</sup>E<sub>g</sub>) transitions, respectively. Because of the low intensity of <sup>2</sup>B<sub>1g</sub> → <sup>2</sup>B<sub>2g</sub>, this band is usually not observed separately in the tetragonally distorted complexes<sup>9</sup>. Electronic spectral data suggesting that an appreciable distortion from octahedral geometry in Cu (II) complexes<sup>9</sup> (Figure 1). All the Ru (II) complexes exhibit four transitions corresponding to <sup>1</sup>A<sub>1g</sub> → <sup>3</sup>T<sub>1g</sub>; <sup>1</sup>A<sub>1g</sub> → <sup>3</sup>T<sub>2g</sub>; <sup>1</sup>A<sub>1g</sub> → <sup>1</sup>T<sub>1g</sub> and <sup>1</sup>A<sub>1g</sub> → <sup>1</sup>T<sub>2g</sub>. Two bands are observed in the 200–530 nm region at the longer wavelength (460–520 nm) have been assigned to the spin allowed <sup>1</sup>A<sub>1g</sub> → <sup>1</sup>T<sub>1g</sub> transition based on molar extinction coefficients. The other high intensity band, at 260–300 nm region has been assigned to the charge-transfer transition arising from the excitation of an electron from the metal t<sub>2g</sub> level to the unfilled molecular orbital derived from the Π\* level of the ligands, in accordance with the assignments confirms octahedral geometry for Ru(II) complexes<sup>25</sup> (Figure 1). The electronic spectra of Pd (II) complexes show a broad d–d transition band in the region of 495–450 nm assignable to <sup>1</sup>B<sub>1g</sub> ← <sup>1</sup>A<sub>1g</sub> transition<sup>23</sup>. Further, a relatively strong charge transfer band has been observed in the spectra of all the Pd (II) complexes in the range of 320–280 nm. From the electronic spectroscopic data and the diamagnetic behavior of the complexes, the square planar geometry has been proposed to all the Pd (II) complexes (Figure 1).



**Figure 1: Proposed Structures of all Schiff base Metal Complexes**

### Conductance measurements

The molar conductance values for the Cu (II) and Ru (II) complexes are determined in dichloromethane and these values are found to be low ( $14 - 16 \text{ ohm}^{-1}\text{cm}^2 \text{ mol}^{-1}$ ) indicating non-electrolytic nature. Two Pd (II) complexes were determined in dimethyl formamide. The conductance values were found  $59$  and  $63 \text{ ohm}^{-1}\text{cm}^2 \text{ mol}^{-1}$  indicating 1:2 electrolytic nature. The electrolytic nature of these Pd (II) complexes is due to the presence of two chloride ions outside the coordination sphere. The presence of chloride ions Pd (II) complexes is detected by the addition of silver nitrate reagent leading to the formation of white precipitate of silver chloride<sup>4, 23-25</sup>.

### Mass spectral analysis

In the mass spectra of PMMA ligand molecular ion peak was observed at  $m/z$   $[M+1]$  403 and HBAB ligand gives molecular ion peak at  $m/z$   $[M+1]$  361. The proposed molecular formula of all the metal complexes was confirmed by comparing their molecular formula weights with  $m/z$  values. The molecular ion peaks at  $m/z$  (M+) 494 ( $[\text{Cu}(\text{HBAB})\text{Cl}_2]$ ), 536 ( $[\text{Cu}(\text{PMAA})\text{Cl}_2]$ ), 532 ( $[\text{Ru}(\text{HBAB})\text{Cl}_2]$ ), 574 ( $[\text{Ru}(\text{PMAA})\text{Cl}_2]$ ), 537 ( $[\text{Pd}(\text{HBAB})\text{Cl}_2]$ ), 579 ( $[\text{Pd}(\text{PMAA})\text{Cl}_2]$ ). This data is in good agreement with the respective molecular formulae of metal complexes.

### Conclusion

In this report the non-template synthesis of six new Schiff base metal complexes. The analytical data show the presence of one metal ion per molecule and suggest a mononuclear structure for all the complexes. In Cu (II) and Ru (II) complexes, two ligands coordinate through four nitrogen (N<sub>4</sub>) atoms/ or two nitrogen and two oxygen (N<sub>2</sub>O<sub>2</sub>) atoms and two chloride ions. These compounds were found to have non-electrolytic nature and distorted octahedral structures were assigned to these complexes. The electronic spectral data are in the favor of square planar structure for Pd (II) metal complexes. Cu (II) and Pd (II) metal complexes showed excellent catalytic activity in different types of organic reactions such as reduction of dabigratin and C-N cross coupling reactions. These Schiff base metal complexes can catalyze selectively and shows moderate to high yields.

## Acknowledgements

The author is very thankful to UGC for sanctioning of minor research project and also thankful to staff of the department for their support in completion of work. The author is specially thankful to Prof.V.Ravinder for continuous encouragement and to Management and principal for providing facilities and continuous their support in completion of work. Finally, author is thankful to the members helped directly or indirectly support in completion of wok.

## References

1. Ashok M, Prasad AVSS, Reddy PM, Ravinder V. Synthesis, Spectral studies, Catalytic and Antibacterial activity of Ru(II) Complexes with Coordinated amides. *Journal of Applied Spectroscopy*. 2008;75(6):872-879.
2. Budige G, Puchakayala MR, Kongara SR, Hu A, Vadde R. Synthesis, characterization and biological evaluation of mononuclear Co(II), Ni(II), Cu(II) and Pd(II) complexes with new N2O2 Schiff base ligands. *Chem Pharm Bull (Tokyo)*. 2011;59(2):166-171.
3. Geeta B, Shrivankumar K, Reddy PM, et al. Binuclear cobalt(II), nickel(II), copper(II) and palladium(II) complexes of a new Schiff-base as ligand: Synthesis, structural characterization, and antibacterial activity. *Spectrochimica Acta Part a-Molecular and Biomolecular Spectroscopy*. Nov 2010;77(4):911-915.
4. Krishna ER, Reddy PM, Sarangapani M, et al. Synthesis of N-4 donor macrocyclic Schiff base ligands and their Ru (II), Pd (II), Pt (II) metal complexes for biological studies and catalytic oxidation of didanosine in pharmaceuticals. *Spectrochimica Acta Part a-Molecular and Biomolecular Spectroscopy*. Nov 2012;97:189-196.
5. Prasad A, Reddy PM, Shanker K, Rohini R, Ravinder V. Application of nickel-catalysed reduction and azo dye reactions for the determination of tinidazole. *Coloration Technology*. 2009;125(5):284-287.
6. Reddy MB, Shanker K, Rani PU, Rohini R, Reddy CK, Ravinder V. New cobalt(II) tetraaza macrocyclic complexes: Synthesis, spectral and antimicrobial studies. *Journal of the Indian Chemical Society*. Oct 2007;84(10):971-977.
7. Reddy PM, Ashok M, Rani PU, Ravinder\* V. Synthesis and Spectral Characterization of Mn (II) Complexes with Amide ligands. *Int J Chem Sci.* 2007;5(2):953-959.
8. Reddy PM, Ho Y-P, Shanker K, Rohini R, Ravinder V. Physicochemical and biological characterization of novel macrocycles derived from o-phthalaldehyde. *European Journal of Medicinal Chemistry*. 2009;44(6):2621-2625.
9. Reddy PM, Prasad AVSS, Ravinder V. Synthesis, spectral characterization, catalytic and antibacterial activity of macrocyclic CuII compounds. *Transition Metal Chemistry*. 2007;32(4):507-513.
10. Reddy PM, Prasad AVSS, Reddy CK, Ravinder V. Synthesis of new macrocyclic rhodium(III) compounds and their utility as catalysts for the oxidation of ascorbic acid. *Transition Metal Chemistry*. Mar 2008;33(2):251-258.
11. Reddy PM, Prasad AVSS, Rohini R, Ravinder V. Catalytic reduction of pralidoxime in pharmaceuticals by macrocyclic Ni(II) compounds derived from orthophthalaldehyde. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2008;70(3):704-712.
12. Reddy PM, Prasad AVSS, Shanker K, Ravinder V. Synthesis, spectral studies and antibacterial activity of novel macrocyclic Co(II) compounds. *Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy*. 2007;68(3):1000-1006.
13. Reddy PM, Rohini R, Krishna ER, Hu AR, Ravinder V. Synthesis, Spectral and Antibacterial Studies of Copper(II) Tetraaza Macrocyclic Complexes. *International Journal of Molecular Sciences*. Apr 2012;13(4):4982-4992.
14. Reddy PM, Shanker K, Srinivas V, et al. Hydrolysis of Letrozole catalyzed by macrocyclic Rhodium (I) Schiff-base complexes. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2015;139:43-48.
15. Rohini R, Reddy PM, Shanker K, Hu AR, Ravinder V. Antimicrobial study of newly synthesized 6-substituted indolo[1,2-c]quinazolines. *European Journal of Medicinal Chemistry*. Mar 2010;45(3):1200-1205.



16. Rohini R, Reddy PM, Shanker K, Hu AR, Ravinder V. Synthesis of Some New Mono, Bis-Indolo[1, 2-c]quinazolines: Evaluation of their Antimicrobial Studies. *Journal of the Brazilian Chemical Society*. 2010;21(5):897-904.
17. Rohini R, Reddy PM, Shanker K, Kanthaiiah K, Ravinder V, Hu A. Synthesis of mono, bis-2-(2-arylideneaminophenyl) indole azomethines as potential antimicrobial agents. *Archives of Pharmacal Research*. Jul 2011;34(7):1077-1084.
18. Rohini R, Reddy PM, Shanker K, Ravinder V. New Mono, Bis-2,2-(arylideneaminophenyl)benzimidazoles: Synthesis and Antimicrobial Investigation. *Acta Chimica Slovenica*. 2009;56(4):900-907.
19. Rohini R, Shanker K, Reddy PM, Ho Y-P, Ravinder V. Mono and bis-6-arylbenzimidazo[1,2-c]quinazolines: A new class of antimicrobial agents. *European Journal of Medicinal Chemistry*. 2009;44:3330–3339.
20. Rohini R, Shanker K, Reddy PM, Ravinder V. Synthesis and Antimicrobial Activities of a New Class of 6-Arylbenzimidazo[1,2-c]quinazolines. *Journal of the Brazilian Chemical Society*. 2010;21(1):49-U36.
21. Rohini R, Shanker K, Reddy PM, Sekhar VC, Ravinder V. 6-Substituted Indolo[1,2-c]quinazolines as New Antimicrobial Agents. *Archiv Der Pharmazie*. Sep 2009;342(9):533-540.
22. Shanker K, Ashok M, Reddy PM, Rohini R, Ravinder V. Spectroscopic characterization and antibacterial activities of Mn(III) complexes containing the tetradentate aza Schiff base ligands. *Int J ChemTech Res*. 2009;1(3):777-783.
23. Shanker K, Reddy PM, Rohini R, Ho YP, Ravinder V. Encapsulation of Pd(II) by N4 and N2O2 macrocyclic ligands: their use in catalysis and biology. *Journal of Coordination Chemistry*. 2009;62(18):3040-3049.
24. Shanker K, Reddy PM, Rohini R, Ravinder V. Synthesis and spectral studies of novel biologically active macrocycles derived from O-phthalaldehyde. *Pharm Lett*. 2009;1:97-107.
25. Shanker K, Rohini R, Ravinder V, Reddy RM, Ho YP. Ru(II) complexes of N-4 and N2O2 macrocyclic Schiff base ligands: Their antibacterial and antifungal studies. *Spectrochimica Acta Part a-Molecular and Biomolecular Spectroscopy*. Jul 2009;73(1):205-211.
26. Shanker K, Rohini R, Shrivankumar K, Reddy PM, Ho YP, Ravinder V. Synthesis of tetraaza macrocyclic Pd-II complexes; antibacterial and catalytic studies. *Journal of the Indian Chemical Society*. Feb 2009;86(2):153-161.
27. Barwiolek M, Babinska M, Kozakiewicz A, Wojtczak A, Kaczmarek-Kedziera A, Szlyk E. Structural and spectral studies of silver(I) complexes with new Schiff bases derived from 2-thiopheneethylamine and their application in thin layer deposition by spin and dip coating techniques. *Polyhedron*. 2017;124:12-21.
28. Fakhari AR, Khorrami AR, Naeimi H. Synthesis and analytical application of a novel tetradentate N2O2 Schiff base as a chromogenic reagent for determination of nickel in some natural food samples. *Talanta*. 2005;66(4):813-817.
29. Jia L, Xu J, Zhao X, et al. Synthesis, characterization, and antitumor activity of three ternary dinuclear copper (II) complexes with a reduced Schiff base ligand and diimine coligands in vitro and in vivo. *Journal of Inorganic Biochemistry*. 2016;159:107-119.
30. Tyagi P, Tyagi M, Agrawal S, Chandra S, Ojha H, Pathak M. Synthesis, characterization of 1,2,4-triazole Schiff base derived 3d-metal complexes: Induces cytotoxicity in HepG2, MCF-7 cell line, BSA binding fluorescence and DFT study. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2017;171:246-257.
31. Cozzi PG. Metal-Salen Schiff base complexes in catalysis: practical aspects. *Chemical Society Reviews*. 2004;33(7):410-421.
32. Chen Y, Liu Y, Zhang X, et al. Stereo-regulated methyl methacrylate (MMA) polymerization catalyzed by asymmetric Salen-type Schiff-base Cu(II) complexes. *Inorganic Chemistry Communications*. 2015;53:1-3.
33. Maurya MR, Chandrakar AK, Chand S. Oxidation of phenol, styrene and methyl phenyl sulfide with H2O2 catalysed by dioxovanadium(V) and copper(II) complexes of 2-aminomethylbenzimidazole-based ligand encapsulated in zeolite-Y. *Journal of Molecular Catalysis A: Chemical*. 2007;263(1–2):227-237.

34. Ashok M, Prasad AVSS, Reddy PM, Ravinder V. Ru(III)-catalyzed oxidation of pyridoxine and albuterol in pharmaceuticals. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2009;72(1):204-208.
35. Reddy PM, Shanker K, Rohini R, Sarangapani M, Ravinder V. Substituted tertiary phosphine Ru(II) organometallics: Catalytic utility on the hydrolysis of etofibrate in pharmaceuticals. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2008;70(5):1231-1237.
36. Rani PU, Reddy PM, Sarangapani M, Ravinder V. Synthesis, spectral and catalytic hydrogenation studies of palladium(II) complexes containing substituted tertiary phosphines. *Journal of the Indian Chemical Society*. Feb 2007;84(2):122-129.
37. Chatterjee D. Olefin epoxidation catalyzed by [RuIII(TDL)(tmeda)H<sub>2</sub>O] complexes (TDL = tridentate Schiff-base ligand; tmeda = tetramethylethylenediamine). *Journal of Molecular Catalysis A: Chemical*. 2009;310(1-2):174-179.
38. Liu S, Peng J, Yang H, Bai Y, Li J, Lai G. Highly efficient and convenient asymmetric hydrosilylation of ketones catalyzed with zinc Schiff base complexes. *Tetrahedron*. 2012;68(5):1371-1375.
39. Jiao T, Tu J, Li G, Xu F. Cu-catalyzed asymmetric Henry reaction promoted by chiral camphor Schiff bases. *Journal of Molecular Catalysis A: Chemical*. 2016;416:56-62.
40. Matsunaga S, Shibasaki M. Recent advances in cooperative bimetallic asymmetric catalysis: dinuclear Schiff base complexes. *Chemical Communications*. 2014;50(9):1044-1057.
41. Kuchtanin V, Kleščíková L, Šoral M, et al. Nickel(II) Schiff base complexes: Synthesis, characterization and catalytic activity in Kumada–Corriu cross-coupling reactions. *Polyhedron*. 2016;117:90-96.
42. Reddy PM, Rohini R, Krishna ER, Hu A, Ravinder V. Synthesis, Spectral and Antibacterial Studies of Copper(II) Tetraaza Macrocyclic Complexes. *International Journal of Molecular Sciences*. 2012;13(4):4982-4992.
43. Shakir M, Khanam S, Azam M, Aatif M, Firdaus F. Template synthesis and spectroscopic characterization of 16-membered [N<sub>4</sub>] Schiff-base macrocyclic complexes of Co(II), Ni(II), Cu(II), and Zn(II): in vitro DNA-binding studies. *Journal of Coordination Chemistry*. 2011/09/20 2011;64(18):3158-3168.
44. Chan KE, Edelman ER, Wenger JB, Thadhani RI, Maddux FW. Dabigatran and Rivaroxaban Use in Atrial Fibrillation Patients on Hemodialysis. *Circulation*. 2015.
45. Ask K, Décologne N, Asare N, et al. Distribution of nitroreductive activity toward nilutamide in rat. *Toxicology and Applied Pharmacology*. 2004;201(1):1-9.
46. Hewitt RG. Abacavir Hypersensitivity Reaction. *Clinical Infectious Diseases*. 2002;34(8):1137-1142.

\*\*\*\*\*