



ChemTech

## International Journal of ChemTech Research

CODEN(USA): IJCRGG, ISSN: 0974-4290, ISSN(Online):2455-9555

Vol.15 No.02, pp 40-47, 2022

### Biological Evaluation of Some Synthesized N-Substituted 1,3,4 Thiadiazole derivatives by using In-Vitro Model

S.R. Kane\*, S. K. Mohite, A. R. Dhole, T. D. Dudhgaokar,  
N. V. Desai, A. M. Desai, A. S. Deshmukh, A. B. Deshmukh

Rajarambapu College of Pharmacy, Kasegaon  
Tal-Walwa Dist-Sangli., MH India

Corres. Author : [kanesandeepr@gmail.com](mailto:kanesandeepr@gmail.com)

**Abstract :** 1,3,4-Thiadiazole and its derivatives continue to be of great interest to a large number of researchers. 1,3,4-thiadiazole is one of the most potent heterocyclic containing carbonic anhydrase and antibacterial inhibitor from the natural and synthetic origin. At present many 1,3,4-thiadiazole nucleus containing drugs are available in the market such as acetazolamide, methazolamide, megalol. In the present investigation we have synthesized 1,3,4 thiadiazoles and tested for brian shrimp lethality assay, possibly due to the presence of =N-C-S moiety. All the synthesized compounds were tested for toxicity by using brian shrimp model. The compound 3d, 3k, 3i, 3n having highest cytotoxic effect according standard drug 5-Fluorouracil.

**Key-words :** 1,3,4-Thiadiazole, toxicity study, Brian shrimp, 5-Fluorouracil..

## Introduction:


1,3,4-Thiadiazole and its derivatives continue to be of great interest to a large number of researchers.<sup>(1)</sup> 1,3,4-thiadiazole is one of the most potent heterocyclic containing carbonic anhydrase and antibacterial inhibitor from the natural and synthetic origin. Till date many 1,3,4-thiadiazole nucleus containing drugs are available in the market such as acetazolamide, methazolamide, megalol.<sup>(2)</sup> 1,3,4 thiadiazoles exhibit diverse biological activities possibly due to the presence of =N-C-S moiety. Moreover, some bi-heterocyclic compounds incorporating 1,3,4thiadiazole or 1,2,4triazole ring have been produced as antimicrobial agents. The cyclization of the compounds having thiosemicarbazide structure has shown to be an excellent strategy for the synthesis of 1,2,4 triazole, 1,3,4 thiadiazole, 1,3,4 oxadiazole and or 1,3,4 triazine derivatives.<sup>(3)</sup> Heterocycles bearing a symmetrical triazole or 1,3,4-thiadiazole moieties, represent an interesting class of compounds possessing a wide spectrum of biological activities such as anti-inflammatory, antiviral and antimicrobial properties. It has also been reported that derivatives of 1,2,4-triazole and 1,3,4-thiadiazole condensed nucleus systems exert diverse pharmacological activities such as anti-inflammatory, antitumor, antifungal and antibacterial.<sup>(4)</sup> A large no of thiadiazoles have been patented in the medicine field for the treatment of a wide variety of diseases and some of them have become commercial products compound in medicinal chemistry because of its therapeutic value .it is also known to have unique antibacterial and anti-inflammatory activities.<sup>(5)</sup> The thiazole moiety belongs to an important class of N and S- containing heterocycles and when attached to a thiourea functional group forms the building block for pharmaceutical agents. They exhibit a wide spectrum of pharmaceutical activity.<sup>(6)</sup> Thiadiazole is a very multifaceted moiety that shows a wide range of pharmacological activity. Thiadiazoles are bioisosteres of heterocycles such as oxadiazole, oxazole and benzene. Substitution of thiadiazoles with their bioisosteres increases activity.<sup>(7)</sup> The potency and selectivity of NSAIDs appear to be directly related to their gastric, renal and hepatotoxicity. A number of 1-acylthiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3-thiones were identified as potent anti-inflammatory compounds.<sup>(8)</sup>

A compound's toxicity is measured using two indicators: the median effective concentration ( $EC_{50}$ ) and the median lethal concentration ( $LC_{50}$ ). The concentration-response curve is generated using 24 h of data to obtain the  $EC_{50}$  (teratogenic effects) and  $LC_{50}$  (embryotoxic effects). The ratio of  $LC_{50}$  to  $EC_{50}$  is the corresponding therapeutic index (TI)<sup>9,10</sup>. In pharmaceutical administration, a TI value < 1 is optimal. In the present study thiourea is important sulfur and nitrogen containing compound and they are useful substances in drug research because it has wide range of Pharmacological activity<sup>11</sup>. The azole when attached to Thiourea functional group forms the building block for pharmacological agents. They exhibit a wide spectrum of pharmaceutical activity. In the design of new Thiourea and 1, 3, 4- thiadiazole derivatives, this novel dual inhibitory activity of enzyme pathways holds promise as anti-inflammatory agent with improved efficacy and safety profile.

## Methodology:<sup>12,13,14,15</sup>

Step 1: Microwave synthesis Method:

- a. Synthesis of 5 (phenyl)-1, 3, 4-thiadiazole -2-amine.
- b. Synthesis of substituted benzoyl isothiocyanate (1a-j).
- c. Synthesis of substituted N- {[5-(phenyl)-1, 3, 4-thiadiazole -2-yl] carbamothioyl} derivatives.



#### A. Synthesis of 1,3,4-Thiadiazole derivatives by microwave method: Step 1<sup>st</sup>: - Synthesis of 5-(substituted)-1,3,4-thiadiazole - 2-amine:

Methyl substituted carboxylate was prepared by using 0.01 mole of substituted carboxylic acid in 35ml methanol. Carboxylic acid is esterified with methanol, The heat the reaction in microwave under reflux for 20 to 25 min at 340 watts.by adding few drops of H<sub>2</sub>SO<sub>4</sub> as catalyst. The mixture of above compound (0.01mol) and 2 ml thiosemicarbazide was refluxed for 15 to 20 min at 34 watts. TLC tested reaction after completion and solid precipitate was dried and recrystallize from methanol the solid precipitate was washed, dried and recrystallize from ethanol.

Step 2<sup>nd</sup>: - Synthesis of substituted benzoyl isothiocyanate:

A chloride substituted for benzoyl (10mmol) solution in C<sub>3</sub>H<sub>6</sub>O (50ml) Was introduced in absolute terms acetone (30ml) to ammonium thiocyanate (10mmol). At 340 watts, The blend of reactions was Shot for 15 to 20 min under reflux. TLC tested after the reaction is complete. refrigerated to room temperature, and filtered off the formed precipitate (NH<sub>4</sub>Cl). To ready for the fresh. To the freshly formulated solution of derivative aroyl isothiocyanate.

Step 3<sup>rd</sup>: - Synthesis of N-[[5-(substituted)-1,3,4-thiadiazole-2-yl] carbamothioyl] derivatives:

A solution of 5-(substituted)-1,3,4-Thiadiazole -2-amine (10mmol) in acetone (10ml) was added and the resulting mixture was stirred with refluxed for 15 to 20 min at 340 watts. TLC tested after the reaction is complete. The solid prod was washed with water and purified by washing with ethanol absolute.The physical data of 5-(substituted)-1,3,4-Thiadiazole -2-amine are presented in table and the physical data of N-[[5-(substituted)-1,3,4-thiadiazole-2-yl] carbamothioyl] are presented in table 6.1.2.2 all the compounds are stable and can be stored at room temperature.

**Table no. 1- Physicochemical data of N-[[5-(substituted)-1,3,4-thiadiazole-2-yl] carbamothioyl] by conventional and microwave method (3a-t).**

Sr. no	Compounds	Molecular Formula	Molecular Weight	MP °C	Percentage Yield%		Rf value
					Conventional	Micro-wave	
1	3a	C <sub>17</sub> H <sub>14</sub> N <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	370.44	134- 136	50%	56%	0.84
2	3b	C <sub>17</sub> H <sub>13</sub> ClN <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	404.89	142- 144	61%	64%	0.91
3	3c	C <sub>17</sub> H <sub>13</sub> ClN <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	404.89	138- 140	66%	70%	0.84
4	3d	C <sub>17</sub> H <sub>12</sub> Cl <sub>2</sub> N <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	439.33	166-168	47%	54%	0.86
5	3e	C <sub>18</sub> H <sub>18</sub> N <sub>4</sub> O <sub>3</sub> S <sub>2</sub>	400.47	144- 146	70%	76%	0.94
6	3f	C <sub>17</sub> H <sub>13</sub> FN <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	388.43	136- 138	61%	67%	0.75
7	3g	C <sub>17</sub> H <sub>13</sub> N <sub>5</sub> O <sub>4</sub> S <sub>2</sub>	415.44	112- 114	73%	79%	0.82
8	3h	C <sub>12</sub> H <sub>11</sub> ClN <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	342.82	126- 128	60%	75%	0.84
9	3i	C <sub>15</sub> H <sub>18</sub> N <sub>4</sub> O <sub>2</sub> S <sub>2</sub>	350.45	124- 126	72%	79%	0.84
10	3j	C <sub>11</sub> H <sub>9</sub> Cl <sub>2</sub> N <sub>4</sub> O <sub>2</sub> S <sub>3</sub>	396.31	136- 138	76%	82%	0.89
11	3k	C <sub>16</sub> H <sub>10</sub> N <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	430.41	102- 104	61%	64%	0.94
12	3l	C <sub>16</sub> H <sub>9</sub> ClN <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	464.86	144-146	65%	69%	0.84
13	3m	C <sub>16</sub> H <sub>9</sub> ClN <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	464.86	152- 154	76%	82%	0.76
14	3n	C <sub>16</sub> H <sub>8</sub> Cl <sub>2</sub> N <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	499.30	124-126	79%	85%	0.74
15	3o	C <sub>17</sub> H <sub>12</sub> N <sub>6</sub> O <sub>6</sub> S <sub>2</sub>	460.44	128- 130	69%	76%	0.78
16	3p	C <sub>16</sub> H <sub>9</sub> FN <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	448.40	118- 120	65%	79%	0.83
17	3q	C <sub>16</sub> H <sub>9</sub> N <sub>7</sub> O <sub>7</sub> S <sub>2</sub>	475.41	128-130	72%	78%	0.92
18	3r	C <sub>11</sub> H <sub>7</sub> ClN <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	402.79	196-198	80%	87%	0.82
19	3s	C <sub>14</sub> H <sub>14</sub> N <sub>6</sub> O <sub>5</sub> S <sub>2</sub>	410.42	144-146	78%	87%	0.84
20	3t	C <sub>10</sub> H <sub>5</sub> Cl <sub>2</sub> N <sub>6</sub> O <sub>5</sub> S <sub>3</sub>	456.28	132-134	82%	88%	0.78

**Cytotoxicity Test**<sup>16,17</sup>

Recently, brine shrimp (*Artemia salina*) lethality assay is commonly used to check the cytotoxic effect of bioactive chemicals. Bioassay of brine shrimp deaths is simple, rapid, inexpensive, easily mastered, and requires small amounts of test material (10-20 mg or less). For those kinds of human solid tumors the bioassay has good correlation with cytotoxic activity.

**Serial dilution of synthesized compound**

All test tubes were clean taken and labeled properly. An analytical balance was weighed on the synthesized 10 mg compound. Stock solution was then prepared with 10 mg soluble of synthesized compound. Serially Prepared dilution from stock solution. Four it has been the test tubes labeled as 1-4.

**Hatching brine shrimp**

Measure 3 liters of water using measuring cylinder and pour into the rectangular jar. Weigh about 27 g of table salt by an balance and supplemented with 4.2 mg/l dried yeast add it into the jar containing water. Mix salt in water with help of spatula. Place the tip of an airline air pump in the bottom of the pot by continually maintaining the aeration. At the top water level of the jar, add about 10 g of brine shrimp eggs and mix with water. Switch on a light, Placed within a few inches of the pot. After 20-24 hours the nauplii will hatch. Observe the nauplii. Select the nauplii after 20-24 hours from next. Hatched nauplii from empty egg should be separated and through the shutting the air off and the light off. Pour 10 nauplii it into test tube by using Pasteur pipette.

**Toxicity testing**

Expose the nauplii to various concentrations of the synthesized compound. Count the no. of live and calculate the percentage of death after 24 hours.

**Calculation:**

Number of % dead nauplii calculated by using following formula,


$$\% \text{ Death} = \frac{\text{Number of dead nauplii}}{\text{Number of dead nauplii} + \text{Number of live nauplii}} \times 100$$

**Observation:**

**Fig.no.1: brine shrimp nauplii dead images.**

Table no.2: Brine shrimp toxicity of synthesized compound (3a-t):

Sr. no	Compound	% dead nauplii			
		1mg	100ug/ml	10ug/ml	1ug/ml
1	3a	100	80	50	10
2	3b	100	80	60	50
3	3c	100	70	40	20
4	<b>3d</b>	<b>100</b>	<b>80</b>	<b>60</b>	<b>10</b>
5	3e	90	70	70	20
6	3f	80	60	60	20
7	3g	90	80	50	10
8	3h	70	70	70	30
9	<b>3i</b>	<b>90</b>	<b>70</b>	<b>70</b>	<b>30</b>
10	3j	100	80	50	30
11	<b>3k</b>	<b>80</b>	<b>70</b>	<b>70</b>	<b>10</b>
12	3l	80	70	50	20
13	3m	100	70	50	40
14	<b>3n</b>	<b>100</b>	<b>80</b>	<b>50</b>	<b>30</b>
15	3o	90	50	40	10
16	3p	100	80	70	30
17	3q	80	70	70	40
18	3r	80	70	70	40
19	3s	90	70	70	30
20	3t	90	50	40	10
21	<b>Standard(5FU)</b>	<b>100</b>	<b>70</b>	<b>70</b>	<b>20</b>



Graph No :1

## Result and discussion:

Check sample lethality test in a zoological organism, such as the saline Artemia (shrimp). Brian Shrimp Cytotoxicity Test sampling a large scale compounds for their various bioactivities is very useful. The synthesized compound 3a-t used for the cytotoxicity test by using brine shrimp. The compound 3d, 3k, 3i, 3n having highest cytotoxic effect according standard drug 5-Fluorouracil.

## References:

1. Gabe K ,Five-membered Rings with More than Two Heteroatoms and Fused Carbocyclic Derivatives,Comprehensive Heterocyclic Chemistry II, 1996.
2. Vinit R, Amit R, Mahendra S, Arvind K, Vinod K , Sharma S, Recent updates on 1,3,4 thiadiazoles : As Anticonvulsant activity ,EC Pharmaceutical Science 2.1(2015) 202 – 229 .
3. Demirbas N, Karaoglu SA, Demirbas A, Sancak K, Synthesis and antimicrobial activities of some new 1-(5-phenylamino-[1,3,4]thiadiazol-2-yl)methyl-5-oxo-[1,2,4]triazole and 1-(4-phenyl-5-thioxo-[1,2,4]triazol-3-yl)methyl-5-oxo-[1,2,4]triazole derivatives, Eur. J. Med. Chem.39 (2004) 793–804.
4. Mohammad A, Kumar H, Javed SA, Condensed bridgehead nitrogen heterocyclic system: Synthesis and pharmacological activities of 1,2,4-triazolo-[3,4-b]-1,3,4-thiadiazole derivatives of ibuprofen and biphenyl-4-yloxy acetic acid, Eur. J. Med. Chem. 43 (2007) 2056-2066.
5. Parmar CK, Umrigar HN, Review article on synthesis of 1, 3, 4-thiadiazole derivatives and its biological activity, J. chem. pharm. res. 9 (2017)202-214.
6. Saeed S, Rashid N, Jones GP, Hussain R, Bhatti HM,Synthesis, spectroscopic characterization, crystal structure and antifungal activity of thiourea derivatives containing a thiazole moiety,Cent. Eur. J. Chem.8 (2009) 550-558.
7. Dr. Annapoorna V, 2, 5- Disubstituted 1, 3, 4-Thiadiazoles: Molecules of Manifold Applications – A Review. 8 (2019) 218-240.
8. Erhan P, Gulay S, Pelin K, N TD, Gulcin A, Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3- thiones, Il Farmaco. (2002) 101–107.
9. Yang Hu,Cui-Yun Li,Xiao-Ming Wang, Yong-Hua Yang, Hai-Liang Zhu, 1, 3, 4-Thiadiazole: Synthesis, Reactions and Applications in Medicinal, Agriculturaland Materials Chemistry, Chemical Reviews. (2005) 1-39.
10. Khan I, Ali S , Hameed S , Rama HN , Hussain TM , Wadood A, Uddin R , Ul-Haq Z, Khan A, Ali S , Choudhary IM, Synthesis, antioxidant activities and urease inhibition of some new 1, 2, 4-triazole and 1,3,4-thiadiazole derivatives,Eur. J. Med. Chem. (2010) 5200-5207.
11. Khatoon Y, Shaquiquzzaman M ,Singh V , Sarafroz M, synthesis of some new 2,5 disubstituted – 1,3,4 –thiadiazole derivatives and investigation of their anticonvulsant activities, Int.J.pharm.sci., 43 (2017) 231 – 237.
12. Subbaiah N, Mohanty S, Sudha B, Ayyann C,Synthesis, characterization and evaluation of antibacterial and antifungal activity of 2,5-disubstituted 1,3,4-oxadaizole derivative, International Journal of Research in Pharmacy and Pharmaceutical Science. 1 (2016) 39-42.
13. Patel BM, Modi RN, Raval PJ, Menon KS,Calix [4] arene based 1,3,4-Oxaadizole and thiadiazole derivatives: Design, synthesis, and biological evaluation, Organic and Biomolecular Chemistry. (2012) 1785-1794.
14. Terzioglu N, Gursoy A,Synthesis and anticancer evaluation of some new hydrazone derivatives of 2,6 dimethylimidazo[2,1-b] - [1,3,4]thiadiazole-5-carbohydrazide,European Journal of Medicinal Chemistry. (2003) 781-786.
15. Gebrelibanos Hiben, M.; Kamelia, L.; de Haan, L.; Spengelink, B.; Wesseling, S.; Vervoort, J.; Rietjens, I.M.C.M. Hazard assessment of Maerua subcordata (Gilg) DeWolf. for selected endpoints using a battery of in vitro tests. J. Ethnopharmacol. (2019), 241, 111978.
16. Wibowo, I.; Permadi, K.; Hartati, R.; Damayanti, S. Ethanolic extract of pomegranate (Punica granatum L) peel: Acute toxicity tests on Brian shrimp embryos and its toxicity prediction by in silico. J. Appl. Pharm. Sci. (2018), 8, 082–086.

17. Yamashita, M. Apoptosis in Brian shrimp development. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* (2003), 136, 731–742.

\*\*\*\*\*