

Antibacterial Activity of Extracts from *Hypnea valentiae* against standard pathogenic Bacterial Strains

Prasanna Latha Dunga* and K.P.J. Hema Latha.

Department of Microbiology,
Andhra University, Visakhapatnam (A.P) – 530 003, India

*Corres Author: prasanna.phd@rediffmail.com
Mobile number: 9949687827

Abstract: The principal objective of the present research work was to determine the antibacterial potential of *Hypnea valentiae* against seven standard pathogenic bacterial strains. To evaluate antibacterial activity the agar-well diffusion assay was used. All the three extracts showed the highest and significant antibacterial activity against both Gram-negative and Gram-positive bacteria. It is the strain *Bacillus subtilis* that is almost resistant to the three extracts of *Hypnea valentiae*.

Key words: Antibacterial activity, Standard pathogenic bacteria, *Hypnea valentiae*.

Introduction:

Use of traditional medicine is one of the common practices in India due to their wide, pharmacological activities¹. Developed and developing countries use traditional medicine at the primary health care level. Many currently used drugs are expensive or not readily available and a major set back to their continued usage is the development of resistance. This situation urgently forced scientists for searching new, inexpensive drugs that will be able to act for longer periods before resistance sets in².

The main aim of the present research work was to determine the antibacterial potential of *Hypnea valentiae* against seven standard pathogenic bacterial strains. This selection was guided by lack or incompleteness of information in literature on antibacterial activity of their extracts; Three extracts of *Hypnea valentiae* were tested against both Gram negative (*Escherichia coli* ATCC 9637, *Klebsiella pneumoniae* MTCC 2405, *Proteus vulgaris* MTCC

0426 and *Micrococcus luteus* MTCC 0439) and Gram-positive (*Bacillus subtilis* MTCC 2274, *Enterococcus faecalis* MTCC 0439 and *Streptococcus faecalis* MTCC 0459) bacteria.³⁻⁵

Experimental

Algal material:

The algae (Red algae) *Hypnea valentiae* was collected from the Tenneti Park, beach, Visakhapatnam, A.P. The algae was identified in the Department of Botany, Andhra University by comparison with herbarium specimen of the Department. The algae was dried under shade for 3 weeks and pulverized with a mechanical grinder.

Preparation of Extracts:

Different extracts were prepared by a modification of the method according to Robinson³; 140g of algal powder was macerated with ethanol in a conical flask for 24 hrs. Mother liquor (crude ethanol extract) was filtered out and the dried

residual algal material was again macerated with hexane (60-80°C) for 24 hrs, filtered and was evaporated to dryness at room temperature (hexane extract). The residual algal material after macerated with hexane was again macerated with methanol (cold) for 48 hrs, followed and was evaporated to obtain cold methanol extract. The residual algal material after cold maceration with methanol was again subjected to soxhlet apparatus and exhaustively extracted with methanol (hot) until solvent turned pure and colourless. The methanol extract was filtered and evaporated under reduced pressure using rotavapor (Heidolph, Heizbad, Laborota 4001, Germany, 2000). All extracts were stored in sterilized containers at room temperature until used for antibacterial testing. At the time of testing, the extracts were reconstituted to a concentration of 100 mg/ml in Dimethyl sulphoxide (DMSO).

Screening of Antibacterial Assay

The antibacterial activity was tested by agar well diffusion method⁴, Mueller–Hinton broth was applied for growing and diluting the bacterial suspensions. Bacterial strains were grown to exponential phase in Mueller – Hinton at 37°C for 18 hrs and adjusted to a final density of 10⁸ CFU/ml by diluting fresh cultures and comparison with Mc Farland density.

For susceptibility testing, 500 ml of Mueller – Hinton broth was inoculated into 250ml of nutrient agar and allowed them to cool under strict aseptic conditions. After the medium was solidified a well was made in petriplates with the help of a sterite metal borer (6mm). 50µl of each extracts were filled in each well by using adjustable volume digital finn pipette. After that the plates were incubated at 37°C for 24 hrs. After proper incubation, antibacterial activity was determined by measuring the diameter of the zone of the inhibition around the well by using Hiantibiotic zone scale – C and the activity was compared with Ciprofloxacin (5µg/ml) (Multidisk, Axiom, Axiom Laboratories, New Delhi). Simultaneously, control (DMSO) was also maintained with extract. There replicates were

carried out for each extract against each of the test organism.

Results and Discussion

The antibacterial activity of crude extracts (hexane, cold methanol and hot methanol extracts at a concentration of 100 mg/ml) showing positive results; The results of diameters of the zones of inhibition of extracts and antibiotic are presented in Table-1, and were interpreted as sensitive (18mm), intermediate (14-17 mm) and resistant⁵ (< 14 mm).

Hexane extract of *Hypnea Valentiae* was found effective against *Klebsiella pneumoniae* (17 mm), *Micrococcus luteus* (14 mm), *Enterococcus faecalis* (18 mm) and *Streptococcus faecalis* (19 mm). Bacterial species *Escherichia coli* and *Proteus vulgaris* were found to be resistant (IZ = 11-13mm). Cold methanol extract showed promising result against all tested bacteria (IZ = range between 18-27 mm) except *Klebsiella pneumoniae* and *Micrococcus luteus* were found to be resistant (IZ = 12 mm). Hot methanol extract showed high antibacterial activity against all tested Gram – negative and Gram – positive bacteria (IZ = range between 14-29 mm). Where as three extracts of *Hypnea valentiae* did not show any antibacterial activity against *Bacillus subtilis*. Most of the bacterial species showed fairly high degree of sensitivity to the methanol (both cold and hot) extracts of *Hypnea valentiae* (IZ = ranging between 17-29mm). It indicates *Hypnea valentiae* may serve as source for compounds with therapeutic potency.

Although this study investigated the preliminary screening of antibacterial activity, the results showed that the extracts from *Hypnea valentiae* possess good antibacterial activity, conferring the great potentials of this algae used in folk-lore medicine for the production of bioactive compounds and are useful for rationalizing the use of this Red algae in preliminary health care. It is therefore, the above findings recommended the further investigation on isolation and purification of bioactive compound responsible for the antibacterial activity.

Table-1 Antibacterial Activity of Extracts from *Hypnea valentiae* against standard Bacterial strains.

Extracts	Test Organisms*						
	E.c	K.n	P.v	M...	B.s	E.f	S.f
Hexane	13	17	11	14	--	18	19
Cold methanol	18	12	18	12	--	20	27
Hot methanol	19	14	20	16	--	21	29
Ciprofloxacin	15	20	20	22	21	14	22

* E.c: *Escherichia coli*, K.p : *Klebsiella pneumoniae*

P.v: *Proteus vulgaris*, M....: *Micrococcus luteus*,

B.s: *Bacillus subtilis*, E.f: *Enterococcus faecalis*,

S.f: *Streptococcus faecalis*. Values are the mean of three replicates; -- : no inhibition.

Acknowledgements

The Authors are thankful to the Department of Microbiology, Andhra University, Visakhapatnam for all the necessary help that was provided to the study.

References:

1. Reichelt JL, Borowitzka MA. Antimicrobial activity from marine algae: results of a large-scale screening program, *Hydrobiologia* 116/117, 1984, 158 ± 167.
2. Del Val AG, Platas G, Basilio A, gorrochategui J, Suai I, Vicente F, Portilo E, del Rio MJ, Reina GG and Pelaez, F Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *Int Microbiol.*, 4, 2001, 35-40.
3. Robinson, T., The organic constituent of higher plants. Their chemistry and interrelationships. Burgess Publishing Company, USA, 1963, 306pp.
4. Karaman, I., Sahin, F., Gulluce, M., Oguteu, H; Sengul, M. and Adiguel, A., Antimicrobial activity of aqueous and methanol extracts of *Juniperus oxycedrus* L., *J. Ethnopharmacol.*, 2003, 85, 231-235.
5. Barry, A.J.M., Garcia, F. and Thrupp, L.D., Interpretation of sensitivity test results, *American Journal of Clinical Pathology*, 1970, 53, 140.
